

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(1) International Patent Classification <sup>5</sup> : A61K 37/12, C07K 3/00, 13/00 C07K 15/00, 17/00		A1	(11) International Publication Number: WO 92/15322
			(43) International Publication Date: 17 September 1992 (17.09.92)
(21) International Application Number: PCT/US92/01785		(74) Agent: CLARK, Paul, T.; Fish & Richardson, 225 Franklin Street, Boston, MA 02110-2804 (US).	
(22) International Filing Date: 6 March 1992 (06.03.92)			
(30) Priority data: 665,961 7 March 1991 (07.03.91) US		(81) Designated States: AT (European patent), AU, BE (European patent), BR, CA, CH (European patent), CS, DE (European patent), DK (European patent), ES (European patent), FI, FR (European patent), GB (European patent), GR (European patent), HU, IT (European patent), JP, KR, LU (European patent), MC (European patent), NL (European patent), NO, PL, RU, SE (European patent).	
(71) Applicant: THE GENERAL HOSPITAL CORPORATION [US/US]; Office of Technology Affairs, Thirteenth Street, Building 149, Suite 1101, Charlestown, MA 02129 (US).			

références, formules, pages à photocopier, etc	No	classement
	1	C07K 14/705
	2	C07K 15/00
	3	C07K 17/00
	4	C07K 18/00
	5	C07K 19/00
	6	C07K 20/00
	7	C07K 21/00
	8	C07K 22/00
	9	C07K 23/00
	10	C07K 24/00
	11	C07K 25/00
	12	C07K 26/00
	13	C07K 27/00
	14	C07K 28/00
	15	C07K 29/00
	16	C07K 30/00
	17	C07K 31/00
	18	C07K 32/00
	19	C07K 33/00
	20	C07K 34/00
	21	C07K 35/00
	22	C07K 36/00
	23	C07K 37/00
	24	C07K 38/00
	25	C07K 39/00
	26	C07K 40/00
	27	C07K 41/00
	28	C07K 42/00
	29	C07K 43/00
	30	C07K 44/00
	31	C07K 45/00
	32	C07K 46/00
	33	C07K 47/00
	34	C07K 48/00
	35	C07K 49/00
	36	C07K 50/00
	37	C07K 51/00
	38	C07K 52/00
	39	C07K 53/00
	40	C07K 54/00
	41	C07K 55/00
	42	C07K 56/00
	43	C07K 57/00
	44	C07K 58/00
	45	C07K 59/00
	46	C07K 60/00
	47	C07K 61/00
	48	C07K 62/00
	49	C07K 63/00
	50	C07K 64/00
	51	C07K 65/00
	52	C07K 66/00
	53	C07K 67/00
	54	C07K 68/00
	55	C07K 69/00
	56	C07K 70/00
	57	C07K 71/00
	58	C07K 72/00
	59	C07K 73/00
	60	C07K 74/00
	61	C07K 75/00
	62	C07K 76/00
	63	C07K 77/00
	64	C07K 78/00
	65	C07K 79/00
	66	C07K 80/00
	67	C07K 81/00
	68	C07K 82/00
	69	C07K 83/00
	70	C07K 84/00
	71	C07K 85/00
	72	C07K 86/00
	73	C07K 87/00
	74	C07K 88/00
	75	C07K 89/00
	76	C07K 90/00
	77	C07K 91/00
	78	C07K 92/00
	79	C07K 93/00
	80	C07K 94/00
	81	C07K 95/00
	82	C07K 96/00
	83	C07K 97/00
	84	C07K 98/00
	85	C07K 99/00
	86	C07K 100/00
	87	C07K 101/00
	88	C07K 102/00
	89	C07K 103/00
	90	C07K 104/00
	91	C07K 105/00
	92	C07K 106/00
	93	C07K 107/00
	94	C07K 108/00
	95	C07K 109/00
	96	C07K 110/00
	97	C07K 111/00
	98	C07K 112/00
	99	C07K 113/00
	100	C07K 114/00
	101	C07K 115/00
	102	C07K 116/00
	103	C07K 117/00
	104	C07K 118/00
	105	C07K 119/00
	106	C07K 120/00
	107	C07K 121/00
	108	C07K 122/00
	109	C07K 123/00
	110	C07K 124/00
	111	C07K 125/00
	112	C07K 126/00
	113	C07K 127/00
	114	C07K 128/00
	115	C07K 129/00
	116	C07K 130/00
	117	C07K 131/00
	118	C07K 132/00
	119	C07K 133/00
	120	C07K 134/00
	121	C07K 135/00
	122	C07K 136/00
	123	C07K 137/00
	124	C07K 138/00
	125	C07K 139/00
	126	C07K 140/00
	127	C07K 141/00
	128	C07K 142/00
	129	C07K 143/00
	130	C07K 144/00
	131	C07K 145/00
	132	C07K 146/00
	133	C07K 147/00
	134	C07K 148/00
	135	C07K 149/00
	136	C07K 150/00
	137	C07K 151/00
	138	C07K 152/00
	139	C07K 153/00
	140	C07K 154/00
	141	C07K 155/00
	142	C07K 156/00
	143	C07K 157/00
	144	C07K 158/00
	145	C07K 159/00
	146	C07K 160/00
	147	C07K 161/00
	148	C07K 162/00
	149	C07K 163/00
	150	C07K 164/00
	151	C07K 165/00
	152	C07K 166/00
	153	C07K 167/00
	154	C07K 168/00
	155	C07K 169/00
	156	C07K 170/00
	157	C07K 171/00
	158	C07K 172/00
	159	C07K 173/00
	160	C07K 174/00
	161	C07K 175/00
	162	C07K 176/00
	163	C07K 177/00
	164	C07K 178/00
	165	C07K 179/00
	166	C07K 180/00
	167	C07K 181/00
	168	C07K 182/00
	169	C07K 183/00
	170	C07K 184/00
	171	C07K 185/00
	172	C07K 186/00
	173	C07K 187/00
	174	C07K 188/00
	175	C07K 189/00
	176	C07K 190/00
	177	C07K 191/00
	178	C07K 192/00
	179	C07K 193/00
	180	C07K 194/00
	181	C07K 195/00
	182	C07K 196/00
	183	C07K 197/00
	184	C07K 198/00
	185	C07K 199/00
	186	C07K 200/00
	187	C07K 201/00
	188	C07K 202/00
	189	C07K 203/00
	190	C07K 204/00
	191	C07K 205/00
	192	C07K 206/00
	193	C07K 207/00
	194	C07K 208/00
	195	C07K 209/00
	196	C07K 210/00
	197	C07K 211/00
	198	C07K 212/00
	199	C07K 213/00
	200	C07K 214/00
	201	C07K 215/00
	202	C07K 216/00
	203	C07K 217/00
	204	C07K 218/00
	205	C07K 219/00
	206	C07K 220/00
	207	C07K 221/00
	208	C07K 222/00
	209	C07K 223/00
	210	C07K 224/00
	211	C07K 225/00
	212	C07K 226/00
	213	C07K 227/00
	214	C07K 228/00
	215	C07K 229/00
	216	C07K 230/00
	217	C07K 231/00
	218	C07K 232/00
	219	C07K 233/00
	220	C07K 234/00
	221	C07K 235/00
	222	C07K 236/00
	223	C07K 237/00
	224	C07K 238/00
	225	C07K 239/00
	226	C07K 240/00
	227	C07K 241/00
	228	C07K 242/00
	229	C07K 243/00
	230	C07K 244/00
	231	C07K 245/00
	232	C07K 246/00
	233	C07K 247/00
	234	C07K 248/00
	235	C07K 249/00
	236	C07K 250/00
	237	C07K 251/00
	238	C07K 252/00
	239	C07K 253/00
	240	C07K 254/00
	241	C07K 255/00
	242	C07K 256/00
	243	C07K 257/00
	244	C07K 258/00
	245	C07K 259/00
	246	C07K 260/00
	247	C07K 261/00
	248	C07K 262/00
	249	C07K 263/00
	250	C07K 264/00
	251	C07K 265/00
	252	C07K 266/00
	253	C07K 267/00
	254	C07K 268/00
	255	C07K 269/00
	256	C07K 270/00
	257	C07K 271/00
	258	C07K 272/00
	259	C07K 273/00
	260	C07K 274/00
	261	C07K 275/00
	262	C07K 276/00
	263	C07K 277/00
	264	C07K 278/00
	265	C07K 279/00
	266	C07K 280/00
	267	C07K 281/00
	268	C07K 282/00
	269	C07K 283/00
	270	C07K 284/00
	271	C07K 285/00
	272	C07K 286/00
	273	C07K 287/00
	274	C07K 288/00
	275	C07K 289/00
	276	C07K 290/00
	277	C07K 291/00
	278	C07K 292/00
	279	C07K 293/00
	280	C07K 294/00
	281	C07K 295/00
	282	C07K 296/00
	283	C07K 297/00
	284	C07K 298/00
	285	C07K 299/00
	286	C07K 300/00
	287	C07K 301/00
	288	C07K 302/00
	289	C07K 303/00
	290	C07K 304/00
	291	C07K 305/00
	292	C07K 306/00
	293	C07K 307/00
	294	C07K 308/00
	295	C07K 309/00
	296	C07K 310/00
	297	C07K 311/00
	298	C07K 312/00
	299	C07K 313/00
	300	C07K 314/00
	301	C07K 315/00
	302	C07K 316/00
	303	C07K 317/00
	304	C07K 318/00
	305	C07K 319/00
	306	C07K 320/00
	307	C07K 321/00
	308	C07K 322/00
	309	C07K 323/00
	310	C07K 324/00
	311	C07K 325/00
	312	C07K 326/00
	313	C07K 327/00
	314	C07K 328/00
	315	C07K 329/00
	316	C07K 330/00
	317	C07K 331/00
	318	C07K 332/00
	319	C07K 333/00
	320	C07K 334/00
	321	C07K 335/00
	322	C07K 336/00
	323	C07K 337/00
	324	C07K 338/00
	325	C07K 339/00
	326	C07K 340/00
	327	C07K 341/00
	328	C07K 342/00
	329	C07K 343/00
	330	C07K 344/00
	331	C07K 345/00
	332	C07K 346/00
	333	C07K 347/00
	334	C07K 348/00
	335	C07K 349/00
	336	C07K 350/00
	337	C07K 351/00
	338	C07K 352/00
	339	C07K 353/00
	340	C07K 354/00
	341	C07K 355/00
	342	C07K 356/00
	343	C07K 357/00
	344	C07K 358/00
	345	C07K 359/00
	346	C07K 360/00
	347	C07K 361/00
	348	C07K 362/00
	349	C07K 363/00
	350	C07K 364/00
	351	C07K 365/00
	352	C07K 366/00
	353	C07K 367/00
	354	C07K 368/00
	355	C07K 369/00
	356	C07K 370/00
	357	C07K



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>5</sup> : A61K 37/12, C07K 3/00, 13/00 C07K 15/00, 17/00	A1	(11) International Publication Number: WO 92/15322 (43) International Publication Date: 17 September 1992 (17.09.92)
--	----	---

(21) International Application Number: PCT/US92/01785

(22) International Filing Date: 6 March 1992 (06.03.92)

(30) Priority data:  
665,961 7 March 1991 (07.03.91) US

(71) Applicant: THE GENERAL HOSPITAL CORPORATION [US/US]; Office of Technology Affairs, Thirteenth Street, Building 149, Suite 1101, Charlestown, MA 02129 (US).

(72) Inventors: SEED, Brian ; 47A Joy Street, Boston, MA 02114 (US). ROMEO, Charles ; 45 Chester Road, Belmont, MA 02178 (US). KOLANUS, Waldemar ; 81 Templeton Parkway, Watertown, MA 02172 (US).

(74) Agent: CLARK, Paul, T.; Fish &amp; Richardson, 225 Franklin Street, Boston, MA 02110-2804 (US).

(81) Designated States: AT (European patent), AU, BE (European patent), BR, CA, CH (European patent), CS, DE (European patent), DK (European patent), ES (European patent), FI, FR (European patent), GB (European patent), GR (European patent), HU, IT (European patent), JP, KR, LU (European patent), MC (European patent), NL (European patent), NO, PL, RU, SE (European patent).

Published  
With international search report.

(54) Title: REDIRECTION OF CELLULAR IMMUNITY BY RECEPTOR CHIMERAS

## (57) Abstract

Disclosed is a method of directing a cellular response in a mammal by expressing in a cell of the mammal a chimeric receptor which causes the cells to specifically recognize and destroy an infective agent, a cell infected with an infective agent, a tumor or cancerous cell, or an autoimmune-generated cell. Also disclosed are cells which express the chimeric receptors and DNA encoding the chimeric receptors.

- 1 -

REDIRECTION OF CELLULAR IMMUNITY BY RECEPTOR CHIMERASField of the Invention

The invention concerns functional T cell receptor, Fc receptor, or B cell receptor chimeras which are capable of redirecting immune system function. More particularly, it concerns the regulation of lymphocytes, macrophages, natural killer cells or granulocytes by the expression in said cells of chimeras which cause the cells to respond to targets recognized by the chimeras. The invention also concerns functional T cell receptor, Fc receptor, or B cell receptor chimeras which are capable of directing therapeutic cells to specifically recognize and destroy either cells infected with a specific infective agent, the infective agent itself, a tumor cell, or an autoimmune-generated cell. More particularly, the invention relates to the production of T cell receptor or Fc receptor chimeras capable of directing cytotoxic T lymphocytes to specifically recognize and lyse cells expressing HIV envelope proteins. The invention therefore provides a therapy for diseases such as AIDS (Acquired Immunodeficiency Syndrome) which are caused by the HIV virus.

Background of the Invention

T cell recognition of antigen through the T cell receptor is the basis of a range of immunological phenomena. The T cells direct what is called cell-mediated immunity. This involves the destruction by cells of the immune system of foreign tissues or infected cells. A variety of T cells exist, including "helper" and "suppressor" cells, which modulate the immune response, and cytotoxic (or "killer") cells, which can kill abnormal cells directly.

A T cell that recognizes and binds a unique antigen displayed on the surface of another cell becomes

- 2 -

activated; it can then multiply, and if it is a cytotoxic cell, it can kill the bound cell.

Autoimmune disease is characterized by production of either antibodies that react with host tissue or  
5 immune effector T cells that are autoreactive. In some instances, autoantibodies may arise by a normal T- and B-cell response activated by foreign substances or organisms that contain antigens that cross react with similar compounds in body tissues. Examples of  
10 clinically relevant autoantibodies are antibodies against acetylcholine receptors in myasthenia gravis; and anti-DNA, anti-erythrocyte, and anti-platelet antibodies in systemic lupus erythematosus.

#### HIV and Immunopathogenesis

15 In 1984 HIV was shown to be the etiologic agent of AIDS. Since that time the definition of AIDS has been revised a number of times with regard to what criteria should be included in the diagnosis. However, despite the fluctuation in diagnostic parameters, the simple  
20 common denominator of AIDS is the infection with HIV and subsequent development of persistent constitutional symptoms and AIDS defining diseases such as a secondary infections, neoplasms, and neurologic disease.  
Harrison's Principles of Internal Medicine, 12th ed.,  
25 McGraw Hill (1991).

HIV is a human retrovirus of the lentivirus group. The four recognized human retroviruses belong to two distinct groups: the human T lymphotropic (or leukemia) retroviruses, HTLV-1 and HTLV-2, and the human  
30 immunodeficiency viruses, HIV-1 and HIV-2. The former are transforming viruses whereas the latter are cytopathic viruses.

HIV-1 has been identified as the most common cause of AIDS throughout the world. Sequence homology between



- 3 -

HIV-2 and HIV-1 is about 40% with HIV-2 being more closely related to some members of a group of simian immunodeficiency viruses (SIV). See Curran, J. et al., Science, 329:1357-1359 (1985); Weiss, R. et al., Nature, 324:572-575 (1986).

HIV has the usual retroviral genes (env, gag, and pol) as well as six extra genes involved in the replication and other biologic activities of the virus. As stated previously, the common denominator of AIDS is a profound immunosuppression, predominantly of cell-mediated immunity. This immune suppression leads to a variety of opportunistic diseases, particularly certain infections and neoplasms.

The main cause of the immune defect in AIDS, has been identified as a quantitative and qualitative deficiency in the subset of thymus-derived (T) lymphocytes, the T4 population. This subset of cells is defined phenotypically by the presence of the CD4 surface molecule, which has been demonstrated to be the cellular receptor for HIV. Dalglish et al., Nature, 312:763 (1984). Although the T4 cell is the major cell type infected with HIV, essentially any human cell that expresses the CD4 molecule on its surface is capable of binding to and being infected with HIV.

Traditionally, CD4<sup>+</sup> T cells have been assigned the role of helper/inducer, indicating their function in providing an activating signal to B cells, or inducing T lymphocytes bearing the reciprocal CD8 marker to become cytotoxic/suppressor cells. Reinherz and Schlossman, Cell, 19:821-827 (1980); Goldstein et al., Immunol. Rev., 68:5-42, (1982).

HIV binds specifically and with high affinity, via a stretch of amino acids in the viral envelope (gp120), to a portion of the V1 region of the CD4 molecule located near its N-terminus. Following binding, the virus fuses

- 4 -

with the target cell membrane and is internalized. Once internalized it uses the enzyme reverse transcriptase to transcribe its genomic RNA to DNA, which is integrated into the cellular DNA where it exists for the life of the cell as a "provirus."

The provirus may remain latent or be activated to transcribe mRNA and genomic RNA, leading to protein synthesis, assembly, new virion formation, and budding of virus from the cell surface. Although the precise mechanism by which the virus induces cell death has not been established, it is felt that the major mechanism is massive viral budding from the cell surface, leading to disruption of the plasma membrane and resulting osmotic disequilibrium.

During the course of the infection, the host organism develops antibodies against viral proteins, including the major envelope glycoproteins gp120 and gp41. Despite this humoral immunity, the disease progresses, resulting in a lethal immunosuppression characterized by multiple opportunistic infections, parasitemia, dementia and death. The failure of the host anti-viral antibodies to arrest the progression of the disease represents one of the most vexing and alarming aspects of the infection, and augurs poorly for vaccination efforts based upon conventional approaches.

Two factors may play a role in the efficacy of the humoral response to immunodeficiency viruses. First, like other RNA viruses (and like retroviruses in particular), the immunodeficiency viruses show a high mutation rate in response to host immune surveillance. Second, the envelope glycoproteins themselves are heavily glycosylated molecules presenting few epitopes suitable for high affinity antibody binding. The poorly antigenic target which the viral envelope presents, allows the host

- 5 -

little opportunity for restricting viral infection by specific antibody production.

Cells infected by the HIV virus express the gp120 glycoprotein on their surface. Gp120 mediates fusion events among CD4<sup>+</sup> cells via a reaction similar to that by which the virus enters the uninfected cells, leading to the formation of short-lived multinucleated giant cells. Syncytium formation is dependent on a direct interaction of the gp120 envelope glycoprotein with the CD4 protein. Dalglish et al., supra; Klatzman, D. et al., Nature, 312:763 (1984); McDougal, J.S. et al., Science, 231:382 (1986); Sodroski, J. et al., Nature, 322:470 (1986); Lifson, J.D. et al., Nature, 323:725 (1986); Sodroski, J. et al., Nature, 321:412 (1986).

Evidence that the CD4-gp120 binding is responsible for viral infection of cells bearing the CD4 antigen includes the finding that a specific complex is formed between gp120 and CD4. McDougal et al., supra. Other investigators have shown that the cell lines, which were noninfective for HIV, were converted to infectable cell lines following transfection and expression of the human CD4 cDNA gene. Maddon et al., Cell, 46:333-348 (1986).

Therapeutic programs based on soluble CD4 as a passive agent to interfere with viral adsorption and syncytium-mediated cellular transmission have been proposed and successfully demonstrated in vitro by a number of groups (Deen et al., Nature, 332:82-84 (1988); Fisher et al., Nature, 331:76-78 (1988); Hussey et al., Nature 331:78-81 (1988); Smith et al., Science, 238:1704-1707 (1987); Trauneker et al., Nature, 331:84-86 (1988)); and CD4 immunoglobulin fusion proteins with extended halflives and modest biological activity have subsequently been developed (Capon et al., Nature, 337:525-531 (1989); Trauneker et al. Nature, 339, 68-70 (1989); Byrn et al., Nature, 344:667-670 (1990);

- 6 -

Zettlmeissl et al., DNA Cell Biol. 9:347-353 (1990)). Although CD4 immunotoxin conjugates or fusion proteins show potent cytotoxicity for infected cells in vitro (Chaudhary et al., Nature, 335:369-372 (1988); Till et al., Science, 242:1166-1168 (1988)), the latency of the immunodeficiency syndrome makes it unlikely that any single-treatment therapy will be effective in eliminating viral burden, and the antigenicity of foreign fusion proteins is likely to limit their acceptability in treatments requiring repetitive dosing. Trials with monkeys affected with SIV have shown that soluble CD4, if administered to animals without marked CD4 cytopenia, can reduce SIV titer and improve in vitro measures of myeloid potential (Watanabe et al., Nature, 337:267-270 (1989)). However a prompt viral reemergence was observed after treatment was discontinued, suggesting that lifelong administration might be necessary to prevent progressive immune system debilitation.

#### T Cell and Fc Receptors

Cell surface expression of the most abundant form of the T cell antigen receptor (TCR) requires the coexpression of at least 6 distinct polypeptide chains (Weiss et al., J. Exp. Med., 160:1284-1299 (1984); Orloffhashi et al., Nature, 316:606-609 (1985); Berkhout et al., J. Biol. Chem., J. Biol. Chem., 263:8528-8536 (1988); Sussman et al., Cell, 52:85-95 (1988)), the  $\alpha/\beta$  antigen binding chains, the three polypeptides of the CD3 complex, and  $\zeta$ . If any of the chains are absent, stable expression of the remaining members of the complex does not ensue.  $\zeta$  is the limiting polypeptide for surface expression of the complete complex (Sussman et al., Cell, 52:85-95 (1988)) and is thought to mediate at least a fraction of the cellular activation programs triggered by receptor recognition of ligand (Weissman et al., EMBO J.,

- 7 -

8:3651-3656 (1989); Frank et al., Science, 249:174-177 (1990)). A 32kDa type I integral membrane homodimer,  $\varsigma$  (zeta) has a 9 residue extracellular domain with no sites for N-linked glycan addition, and a 112 residue (mouse) or 113 residue (human) intracellular domain (Weissman et al., Science, 238:1018-1020 (1988a); Weissman et al., Proc. Natl. Acad. Sci. USA, 85:9709-9713 (1988b)). An isoform of  $\varsigma$  called  $\eta$  (eta) (Baniyash et al., J. Biol. Chem., 263:9874-9878 (1988); Orloff et al., J. Biol. Chem., 264:14812-14817 (1989)), which arises from an alternate mRNA splicing pathway (Jin et al., Proc. Natl. Acad. Sci. USA, 87:3319-3233 (1990)), is present in reduced amounts in cells expressing the antigen receptor.  $\varsigma$ - $\eta$  heterodimers are thought to mediate the formation of inositol phosphates, as well as the receptor-initiated programmed cell death called apoptosis (Merćep et al., Science, 242:571-574 (1988); Merćep et al., Science, 246:1162-1165 (1989)).

Like  $\varsigma$  and  $\eta$ , the Fc receptor-associated  $\gamma$  chain is expressed in cell surface complexes with additional polypeptides, some of which mediate ligand recognition, and others of which have undefined function.  $\gamma$  (gamma) bears a homodimeric structure and overall organization very similar to that of  $\varsigma$ , and is a component of both the mast cell/basophil high affinity IgE receptor, Fc $\epsilon$ RI, which consists of at least three distinct polypeptide chains (Blank et al., Nature, 337:187-189 (1989); Ra et al., Nature, 241:752-754 (1989)), and one of the low affinity receptors for IgG, represented in mice by Fc $\gamma$ RII $\alpha$  (Ra et al., J. Biol. Chem. J. Biol. Chem., 264:15323-15327 (1989)), and in humans by the CD16 subtype expression by macrophages and natural killer cells, CD16<sub>TM</sub> (CD16 transmembrane) (Lanier et al., Nature, 342:803-805 (1989); Anderson et al., Proc. Natl. Acad. Sci. USA, 87:2274-2278 (1990)) and with a polypeptide of

- 8 -

unidentified function (Anderson et al., Proc. Natl. Acad. Sci. USA, 87:2274-2278 (1990)). Recently it has been reported that  $\gamma$  is expressed by a mouse T cell line, CTL, in which it forms homodimers as well as  $\gamma$ - $\zeta$  and  $\gamma$ - $\eta$ .

5 heterodimers (Orloff et al., Nature, 347:189-191 (1990)).

The Fc receptors mediate phagocytosis of immune complexes, transcytosis, and antibody dependent cellular cytotoxicity (ADCC) (Ravetch and Kinet, Annu. Rev. Immunol. 9:457-492 (1991); Unkeless et al., Annu. Rev.

10 Immunol. 6:251-281 (1988); and Mellman, Curr. Opin. Immunol. 1:16-25 (1988)).

Recently it has been shown that one of the murine low affinity Fc receptor isoforms (Fc $\gamma$ RIIB1) mediates internalization of Ig-coated targets into clathrin coated pits, and that another low affinity  
15 receptor (Fc $\gamma$ RIIA) mediates ADCC through its association with one or more members of a small family of 'trigger molecules' (Miettinen et al., Cell 58:317-327 (1989); and Hunziker and Mellman, J. Cell Biol. 109:3291-3302

(1989)). These trigger molecules, T cell receptor (TCR)

20  $\zeta$  chain, TCR  $\eta$  chain, and Fc receptor  $\gamma$  chain, interact with ligand recognition domains of different immune system receptors and can autonomously initiate cellular effector programs, including cytolysis, following aggregation (Samelson et al., Cell 43:223-231 (1985);

25 Weissman et al., Science 239:1018-1020 (1988); Jin et al., Proc. Natl. Acad. Sci. USA 87:3319-3323 (1990);

Blank et al., Nature 337:187-189 (1989); Lanier et al., Nature 342:803-805 (1989); Kurosaki and Ravetch, Nature 342:805-807 (1989); Hibbs et al., Science 246:1608-1611

30 (1989); Anderson et al., Proc. Natl. Acad. Sci. USA 87:2274-2278 (1990); and Irving and Weiss, Cell 64: 891-901 (1991)).

In drawing parallels between the murine and human low affinity Fc receptor families, however, it has become  
35 clear that the human Fc $\gamma$ RIIA and C isoforms have no

- 9 -

murine counterpart. In part because of this, their function has yet to be defined.

Because humoral agents based on CD4 may have limited utility in vivo, the inventors began to explore the possibility of augmenting cellular immunity to HIV. As a result they report the preparation of protein chimeras in which the extracellular domain of CD4 is fused to the transmembrane and/or intracellular domains of T cell receptor, IgG Fc receptor, or B cell receptor signal transducing elements. Cytolytic T cells expressing chimeras which include an extracellular CD4 domain show potent MHC-independent destruction of cellular targets expressing HIV envelope proteins. An extremely important and novel component of this approach has been the identification of single T cell receptor, or Fc receptor, and B cell receptor chains whose aggregation suffices to initiate the cellular response.

One particularly useful application of this approach has been the invention of chimeras between CD4 and  $\zeta$ ,  $\eta$ , or  $\gamma$  that direct cytolytic T lymphocytes to recognize and kill cells expressing HIV gp120.

#### Summary of the Invention

Although native T cell, B cell, and Fc receptors are or can be highly complicated multimeric structures not lending themselves to convenient manipulation, the present invention demonstrates the feasibility of creating chimeras between the intracellular domain of any of a variety of molecules which are capable of fulfilling the task of target recognition. In particular, the formation of chimeras consisting of the intracellular portion of T cell/Fc receptor zeta, eta, or gamma chains joined to the extracellular portion of a suitably engineered antibody molecule allows the target recognition potential of an immune system cell to be

- 10 -

specifically redirected to the antigen recognized by the extracellular antibody portion. Thus with an antibody portion capable of recognizing some determinant on the surface of a pathogen, immune system cells armed with the chimera would respond to the presence of the pathogen with the effector program appropriate to their lineage, e.g., helper T lymphocytes would respond by cytotoxic activity against the target, and B lymphocytes would be activated to synthesize antibody. Macrophages and granulocytes would carry out their effector programs, including cytokine release, phagocytosis, and reactive oxygen generation. Similarly, with an antibody portion capable of recognizing tumor cells, the immune system response to the tumor would be beneficially elevated. With an antibody capable of recognizing immune cells having an inappropriate reactivity with self determinants, the autoreactive cells could be selectively targeted for destruction. Although these examples draw on the use of antibody chimeras as a convenient expository tool, the invention is not limited in scope to antibody chimeras, and indeed, the use of specific nonantibody extracellular domains may have important advantages. For example with an extracellular portion that is the receptor for a virus, bacterium, or parasite, cells armed with the chimeras would specifically target cells expressing the viral, bacterial or parasitic determinants. The advantage of this approach over the use of antibodies is that the native receptor for pathogen may have uniquely high selectivity or affinity for the pathogen, allowing a greater degree of precision in the resulting immune response. Similarly, to delete immune system cells which inappropriately react with a self antigen, it may suffice to join the antigen (either as an intact protein, in the case of B cell depletion therapies, or as MHC complex, in the case of T cell



- 11 -

depletion therapies) to intracellular zeta, eta or gamma chains, and thereby affect the specific targeting of the cells inappropriately responding to self determinants.

Another use of the chimeras is the control of cell  
5 populations in vivo subsequent to other forms of genetic engineering. For example, the use of tumor infiltrating lymphocytes or natural killer cells to carry cytotoxic principles to the site of tumors has been proposed. The present invention provides a convenient means to regulate  
10 the numbers and activity of such lymphocytes and cells without removing them from the body of the patient for amplification in vitro. Thus, because the intracellular domains of the chimeric receptors mediate the proliferative responses of the cells, the coordination of  
15 the extracellular domains by a variety of aggregating stimuli specific for the extracellular domains (e.g., an antibody specific for the extracellular domain) will result in proliferation of the cells bearing the chimeras.

20 Although the specific embodiments of the present invention comprise chimeras between zeta, eta or gamma chains, or active fragments thereof (e.g., those discussed below), any receptor chain having a similar function to these molecules, e.g., in granulocytes or B  
25 lymphocytes, could be used for the purposes disclosed here. The distinguishing features of desirable immune cell trigger molecules comprise the ability to be expressed autonomously (i.e., as a single chain), the ability to be fused to an extracellular domain such that  
30 the resultant chimera is present on the surface of a therapeutic cell, and the ability to initiate cellular effector programs upon aggregation secondary to encounter with a target ligand.

At present the most convenient method for delivery  
35 of the chimeras to immune system cells is through some

- 12 -

form of genetic therapy. However reconstituting immune system cells with chimeric receptors by mixture of the cells with suitably solubilized purified chimeric protein would also result in the formation of an engineered cell population capable of responding to the targets recognized by the extracellular domain of the chimeras. Similar approaches have been used, for example, to introduce the intact HIV receptor, CD4, into erythrocytes for therapeutic purposes. In this case the engineered cell population would not be capable of self renewal.

The present invention relates to functional simplified T cell receptor, B cell receptor, and Fc receptor chimeras which are capable of redirecting immune system function. More particularly, it relates to the regulation of lymphocytes, macrophages, natural killer cells or granulocytes by the expression in said cells of chimeras which cause the cells to respond to targets recognized by the chimeras. The invention also relates to a method of directing cellular response to an infective agent, a tumor or cancerous cell, or an autoimmune generated cell. The method for directing the cellular response in a mammal comprises administering an effective amount of therapeutic cells to said mammal, said cells being capable of recognizing and destroying said infective agent, tumor, cancer cell or autoimmune generated cell.

In another embodiment, the method of directing cellular response to an infective agent comprises administering therapeutic cells capable of recognizing and destroying said agent, wherein the agent is a specific virus, bacteria, protozoa, or fungi. Even more specifically, the method is directed against agents such as HIV and Pneumocystis carinii.

Specifically the invention provides for a method of directing cellular response to an HIV infected cell.

- 13 -

The method comprises administering to a patient an effective amount of cytotoxic T lymphocytes, said lymphocytes being capable of specifically recognizing and lysing cells infected with HIV.

5           Thus, in one embodiment, there is provided according to the invention a method for directing cellular response to HIV infected cells, comprising administering to a patient an effective amount of cytotoxic T lymphocytes which are capable of specifically  
10 recognizing and lysing cells infected with HIV.

          In yet another embodiment is provided the chimeric receptor proteins which direct the cytotoxic T lymphocytes to recognize and lyse the HIV infected cell. Yet another embodiment of the invention comprises host  
15 cells transformed with a vector comprising the chimeric receptors.

          In yet another embodiment, the present invention provides for an antibody against the chimeric receptors of the invention.

20           In order to obtain cytotoxic T lymphocytes which specifically bind and lyse cells infected with HIV, the present inventors therefore attempted, and herein receptor chimeras. These chimeric receptors are functionally active and possess the extraordinary ability  
25 of being able to specifically bind and lyse cells expressing gp120.

          It is an object of the present invention, then, to provide for a method of treatment for individuals infected with HIV. The present invention thus provides a  
30 number of important advances in the therapy of AIDS.

          These and other non-limiting embodiments of the present invention will be apparent to those of skill from the following detailed description of the invention.

          In the following detailed description, reference  
35 will be made to various methodologies known to those of

- 14 -

skill in the art of molecular biology and immunology. Publications and other materials setting forth such known methodologies to which reference is made are incorporated herein by reference in their entireties as though set  
5 forth in full.

Standard reference works setting forth the general principles of recombinant DNA technology include Watson, J.D. et al., Molecular Biology of the Gene, Volumes I and II, the Benjamin/Cummings Publishing Company, Inc.,  
10 publisher, Menlo Park, CA (1987); Darnell, J.E. et al., Molecular Cell Biology, Scientific American Books, Inc., Publisher, New York, N.Y. (1986); Lewin, B.M., Genes II, John Wiley & Sons, publishers, New York, N.Y. (1985); Old, R.W., et al., Principles of Gene Manipulation: An  
15 Introduction to Genetic Engineering, 2d edition, University of California Press, publisher, Berkeley, CA (1981); Maniatis, T., et al., Molecular Cloning: A Laboratory Manual, 2nd Ed. Cold Spring Harbor Laboratory, publisher, Cold Spring Harbor, NY (1989); and Current  
20 Protocols in Molecular Biology, Ausubel et al., Wiley Press, New York, NY (1989).

#### DEFINITIONS

By "cloning" is meant the use of in vitro recombination techniques to insert a particular gene or  
25 other DNA sequence into a vector molecule. In order to successfully clone a desired gene, it is necessary to employ methods for generating DNA fragments for joining the fragments to vector molecules, for introducing the composite DNA molecule into a host cell in which it can  
30 replicate, and for selecting the clone having the target gene from amongst the recipient host cells.

By "cDNA" is meant complementary or copy DNA produced from an RNA template by the action of RNA-dependent DNA polymerase (reverse transcriptase). Thus a

- 15 -

"cDNA clone" means a duplex DNA sequence complementary to an RNA molecule of interest, carried in a cloning vector.

By "cDNA library" is meant a collection of recombinant DNA molecules containing cDNA inserts which  
5 comprise DNA copies of mRNA being expressed by the cell at the time the cDNA library was made. Such a cDNA library may be prepared by methods known to those of skill, and described, for example, in Maniatis et al., Molecular Cloning: A Laboratory Manual, supra.

10 Generally, RNA is first isolated from the cells of an organism from whose genome it is desired to clone a particular gene. Preferred for the purpose of the present invention are mammalian, and particularly human, lymphocytic cell lines. A presently preferred vector for  
15 this purpose is the vaccinia virus WR strain.

By "vector" is meant a DNA molecule, derived, e.g., from a plasmid, bacteriophage, or mammalian or insect virus, into which fragments of DNA may be inserted or cloned. A vector will contain one or more unique  
20 restriction sites and may be capable of autonomous replication in a defined host or vehicle organism such that the cloned sequence is reproducible. Thus, by "DNA expression vector" is meant any autonomous element capable of directing the synthesis of a recombinant  
25 peptide. Such DNA expression vectors include bacterial plasmids and phages and mammalian and insect plasmids and viruses.

By "substantially pure" is meant a compound, e.g., a a protein, a polypeptide, or an antibody, that is  
30 substantially free of the components that naturally accompany it. Generally, a compound is substantially pure when at least 60%, more preferably at least 75%, and most preferably at least 90% of the total material in a sample is the compound of interest. Purity can be  
35 measured by any appropriate method, e.g., column

- 16 -

chromatography, polyacrylamide gel electrophoresis, or HPLC analysis. In the context of a nucleic acid, "substantially pure" means a nucleic acid sequence, segment, or fragment that is free from the genes that flank it in its naturally-occurring state (e.g., free from the sequences that flank the nucleic acid in its native genomic position). By "functional derivative" is meant the "fragments," "variants," "analogues," or "chemical derivatives" of a molecule. A "fragment" of a molecule, such as any of the cDNA sequences of the present invention, is meant to refer to any nucleotide subset of the molecule. A "variant" of such molecule is meant to refer to a naturally occurring molecule substantially similar to either the entire molecule, or a fragment thereof. An "analog" of a molecule is meant to refer to a non-natural molecule substantially similar to either the entire molecule or a fragment thereof. A molecule is said to be "substantially similar" to another molecule if the sequence of amino acids in both molecules is substantially the same. Substantially similar amino acid molecules will possess a similar biological activity. Thus, provided that two molecules possess a similar activity, they are considered variants as that term is used herein even if one of the molecules contains additional or fewer amino acid residues not found in the other, or if the sequence of amino acid residues is not identical. As used herein, a molecule is said to be a "chemical derivative" of another molecule when it contains additional chemical moieties not normally a part of the molecule. Such moieties may improve the molecule's solubility, absorption, biological half life, etc. The moieties may alternatively decrease the toxicity of the molecule, eliminate or attenuate any undesirable side effect of the molecule, etc. Moieties capable of mediating such effects are disclosed, for

- 17 -

example, in Remington's Pharmaceutical Sciences, 16th et., Mack Publishing Co., Easton, Penn. (1980).

Similarly, a "functional derivative" of a receptor chimera of the present invention is meant to include

5 "fragments," "variants," or "analogues" of the gene, which may be "substantially similar" in nucleotide sequence, and which encode a molecule possessing similar activity to, for example, a T cell, B cell, or Fc receptor chimera.

10 Thus, as used herein, a T cell, B cells or Fc receptor chimera protein is also meant to include any functional derivative, fragments, variants, analogues, or chemical derivatives which may be substantially similar to the "wild-type" chimera and which possess similar

15 activity (i.e., most preferably, 90%, more preferably, 70%, preferably 40%, or at least 10% of the wild-type receptor chimera's activity). The activity of a functional chimeric receptor derivative includes specific binding (with its extracellular portion) to a targeted

20 agent or cell and resultant destruction (directed by its intracellular or transmembrane portion) of that agent or cell; such activity may be tested, e.g., using any of the assays described herein.

A DNA sequence encoding the T cell, B cell, or Fc

25 receptor chimera of the present invention, or its functional derivatives, may be recombined with vector DNA in accordance with conventional techniques, including blunt-ended or staggered-ended termini for ligation, restriction enzyme digestion to provide appropriate

30 termini, filling in of cohesive ends as appropriate, alkaline phosphatase treatment to avoid undesirable joining, and ligation with appropriate ligases. Techniques for such manipulations are disclosed by Maniatis, T., et al., supra, and are well known in the

35 art.

- 18 -

A nucleic acid molecule, such as DNA, is said to be "capable of expressing" a polypeptide if it contains nucleotide sequences which contain transcriptional and translational regulatory information and such sequences are "operably linked" to nucleotide sequences which encode the polypeptide. An operable linkage is a linkage in which the regulatory DNA sequences and the DNA sequence sought to be expressed are connected in such a way as to permit gene expression. The precise nature of the regulatory regions needed for gene expression may vary from organism to organism, but shall in general include a promoter region which, in prokaryotes, contains both the promoter (which directs the initiation of RNA transcription) as well as the DNA sequences which, when transcribed into RNA, will signal the initiation of protein synthesis. Such regions will normally include those 5'-non-coding sequences involved with initiation of transcription and translation, such as the TATA box, capping sequence, CAAT sequence, and the like.

If desired, the non-coding region 3' to the gene sequence coding for the protein may be obtained by the above-described methods. This region may be retained for its transcriptional termination regulatory sequences, such as termination and polyadenylation. Thus, by retaining the 3'-region naturally contiguous to the DNA sequence coding for the protein, the transcriptional termination signals may be provided. Where the transcriptional termination signals are not satisfactorily functional in the expression host cell, then a 3' region functional in the host cell may be substituted.

Two DNA sequences (such as a promoter region sequence and a T cell receptor, a B cell receptor, or Fc receptor chimera encoding sequence) are said to be operably linked if the nature of the linkage between the



- 19 -

two DNA sequences does not (1) result in the introduction of a frame-shift mutation, (2) interfere with the ability of the promoter region sequence to direct the transcription of the receptor chimera gene sequence, or  
5 (3) interfere with the ability of the receptor chimera gene sequence to be transcribed by the promoter region sequence. A promoter region would be operably linked to a DNA sequence if the promoter were capable of effecting transcription of that DNA sequence. Thus, to express the  
10 protein, transcriptional and translational signals recognized by an appropriate host are necessary.

The present invention encompasses the expression of a T cell receptor, B cell receptor, or Fc receptor chimera protein (or a functional derivative thereof) in  
15 either prokaryotic or eukaryotic cells, although eukaryotic (and, particularly, human lymphocyte) expression is preferred.

Antibodies according to the present invention may be prepared by any of a variety of methods. For example,  
20 cells expressing the receptor chimera protein, or a functional derivative thereof, can be administered to an animal in order to induce the production of sera containing polyclonal antibodies that are capable of binding the chimera.

25 In a preferred method, antibodies according to the present invention are monoclonal antibodies. Such monoclonal antibodies can be prepared using hybridoma technology (Kohler et al., Nature 256:495 (1975); Kohler et al., Eur. J. Immunol. 6:511 (1976); Kohler et al.,  
30 Eur. J. Immunol. 6:292 (1976); Hammerling et al., In: Monoclonal Antibodies and T-Cell Hybridomas, Elsevier, N.Y., pp. 563-684 (1981)). In general, such procedures involve immunizing an animal with the T cell receptor, B cell receptor, or Fc receptor chimera antigen. The  
35 splenocytes of such animals are extracted and fused with

- 20 -

a suitable myeloma cell line. Any suitable myeloma cell line may be employed in accordance with the present invention. After fusion, the resulting hybridoma cells are selectively maintained in HAT medium, and then cloned  
5 by limiting dilution as described by Wands, J.R., et al. (Gastroenterology 80:225-232 (1981)). The hybridoma cells obtained through such a selection are then assayed to identify clones which secrete antibodies capable of binding the chimera.

10       Antibodies according to the present invention also may be polyclonal, or, preferably, region specific polyclonal antibodies.

Antibodies against the T cell receptor, B cell receptor, or Fc receptor chimera according to the present  
15 invention may be used to monitor the amount of chimeric receptor (or chimeric receptor-bearing cells) in a patient. Such antibodies are well suited for use in standard immunodiagnostic assay known in the art, including such immunometric or "sandwich" assays as the  
20 forward sandwich, reverse sandwich, and simultaneous sandwich assays. The antibodies may be used in any number of combinations as may be determined by those of skill without undue experimentation to effect immunoassays of acceptable specificity, sensitivity, and  
25 accuracy.

Standard reference works setting forth general principles of immunology include Roitt, I., Essential Immunology, Sixth Ed., Blackwell Scientific Publications, Publisher, Oxford (1988); Kimball, J. W., Introduction to  
30 Immunology, Second Ed., Macmillan Publishing Co., Publisher, New York (1986); Roitt, I., et al., Immunology, Gower Medical Publishing Ltd., Publisher, London, (1985); Campbell, A. , "Monoclonal Antibody Technology," in, Burdon, R., et al., eds., Laboratory  
35 Techniques in Biochemistry and Molecular Biology, Volume

- 21 -

- 13, Elsevier, Publisher, Amsterdam (1984); Klein, J.,  
Immunology: The Science of Self-Nonself Discrimination,  
John Wiley & Sons, Publisher, New York (1982); and  
Kennett, R., et al., eds., Monoclonal Antibodies,  
5 Hybridoma: A New Dimension In Biological Analyses,  
Plenum Press, Publisher, New York (1980).

By "detecting" it is intended to include  
determining the presence or absence of a substance or  
quantifying the amount of a substance. The term thus  
10 refers to the use of the materials, compositions, and  
methods of the present invention for qualitative and  
quantitative determinations.

The isolation of other hybridomas secreting  
monoclonal antibodies of the same specificity as those  
15 described herein can be accomplished by the technique of  
anti-idiotypic screening (Potocmjak, et al., Science  
215:1637 (1982)). Briefly, an anti-idiotypic antibody is  
an antibody which recognizes unique determinants present  
on the antibody produced by the clone of interest. The  
20 anti-idiotypic antibody is prepared by immunizing an  
animal of the same strain used as the source of the  
monoclonal antibody with the monoclonal antibody of  
interest. The immunized animal will recognize and  
respond to the idiotypic determinants of the immunizing  
25 antibody by producing antibody to these idiotypic  
determinants (anti-idiotypic antibody).

For replication, the hybrid cells may be  
cultivated both in vitro and in vivo. High in vivo  
production makes this the presently preferred method of  
30 culture. Briefly, cells from the individual hybrid  
strains are injected intraperitoneally into pristane-  
primed BALB/c mice to produce ascites fluid containing  
high concentrations of the desired monoclonal antibodies.  
Monoclonal antibodies of isotype IgM or IgG may be  
35 purified from cultured supernatants using column

- 22 -

chromatography methods well known to those of skill in the art.

Antibodies according to the present invention are particularly suited for use in immunoassays wherein they  
5 may be utilized in liquid phase or bound to a solid phase carrier. In addition, the antibodies in these immunoassays can be detectably labeled in various ways.

There are many different labels and methods of labeling known in the art. Examples of the types of  
10 labels which can be used in the present invention include, but are not limited to, enzymes, radioisotopes, fluorescent compounds, chemiluminescent compounds, bioluminescent compounds and metal chelates. Those of ordinary skill in the art will know of other suitable  
15 labels for binding to antibodies, or will be able to ascertain the same by the use of routine experimentation. Furthermore, the binding of these labels to antibodies can be accomplished using standard techniques commonly known to those of ordinary skill in the art.

20 One of the ways in which antibodies according to the present invention can be detectably labeled is by linking the antibody to an enzyme. This enzyme, in turn, when later exposed to its substrate, will react with the substrate in such a manner as to produce a chemical  
25 moiety which can be detected as, for example, by spectrophotometric or fluorometric means. Examples of enzymes which can be used to detectably label antibodies include malate dehydrogenase, staphylococcal nuclease, delta-V-steroid isomerase, yeast alcohol dehydrogenase,  
30 alpha-glycerophosphate dehydrogenase, triose phosphate isomerase, biotinavidin peroxidase, horseradish peroxidase, alkaline phosphatase, asparaginase, glucose oxidase,  $\beta$ -galactosidase, ribonuclease, urease, catalase, glucose-VI-phosphate dehydrogenase, glucoamylase and  
35 acetylcholine esterase.

- 23 -

The presence of detectably labeled antibodies also can be detected by labeling the antibodies with a radioactive isotope which then can be determined by such means as the use of a gamma counter or a scintillation  
5 counter. Isotopes which are particularly useful for the purpose of the present invention are  $^3\text{H}$ ,  $^{125}\text{I}$ ,  $^{32}\text{P}$ ,  $^{35}\text{S}$ ,  $^{14}\text{C}$ ,  $^{51}\text{Cr}$ ,  $^{36}\text{Cl}$ ,  $^{57}\text{Co}$ ,  $^{58}\text{Co}$ ,  $^{59}\text{Fe}$  and  $^{75}\text{Se}$ .

It is also possible to detect the binding of detectably labeled antibodies by labeling the antibodies  
10 with a fluorescent compound. When a fluorescently labeled antibody is exposed to light of the proper wavelength, its presence then can be detected due to the fluorescence of the dye. Among the most commonly used fluorescent labeling compounds are fluorescein,  
15 isothiocyanate, rhodamine, phycoerythrin, phycocyanin, allophycocyanin, o-phthaldehyde and fluorescamine.

The antibodies of the invention also can be detectably labeled using fluorescent emitting metals such as  $^{152}\text{Eu}$ , or others of the lanthanide series. These  
20 metals can be attached to the antibody molecule using such metal chelating groups as diethylenteraminepentaacetic acid (DTPA) or ethylenediaminetetraacetic acid (EDTA).

Antibodies also can be detectably labeled by  
25 coupling them to a chemiluminescent compound. The presence of the chemiluminescent-tagged antibody is then determined by detecting the presence of luminescence that arises during the course of the chemical reaction. Examples of particularly useful chemiluminescent labeling  
30 compounds are luminal, isoluminol, theromatic acridinium ester, imidazole, acridinium salts, oxalate ester, and dioxetane.

Likewise, a bioluminescent compound may be used to label the antibodies according to the present invention.  
35 Bioluminescence is a type of chemiluminescence found in

- 24 -

biological systems in which a catalytic protein increases the efficiency of the chemiluminescent reaction. The presence of a bioluminescent antibody is determined by detecting the presence of luminescence. Important  
5 bioluminescent compounds for purposes of labeling include luciferin, luciferase aequorin.

The antibodies and substantially purified antigen of the present invention are ideally suited for the preparation of a kit. Such a kit may comprise a carrier  
10 means being compartmentalized to receive in close confinement therewith one or more container means such as vials, tubes and the like, each of said container means comprising the separate elements of the assay to be used.

The types of assays which can be incorporated in  
15 kit form are many, and include, for example, competitive and non-competitive assays. Typical examples of assays which can utilize the antibodies of the invention are radioimmunoassays (RIA), enzyme immunoassays (EIA), enzyme-linked immunosorbent assays (ELISA), and  
20 immunometric, or sandwich, immunoassays.

By the term "immunometric assay" or "sandwich immunoassay," it is meant to include simultaneous sandwich, forward sandwich and reverse sandwich immunoassays. These terms are well understood by those  
25 skilled in the art. Those of skill will also appreciate that antibodies according to the present invention will be useful in other variations and forms of assays which are presently known or which may be developed in the future. These are intended to be included within the  
30 scope of the present invention.

In the preferred mode for performing the assays it is important that certain "blockers" be present in the incubation medium (usually added with the labeled soluble antibody). The "blockers" are added to assure that non-  
35 specific proteins, protease, or human antibodies to mouse

- 25 -

immunoglobulins present in the experimental sample do not cross-link or destroy the antibodies on the solid phase support, or the radiolabeled indicator antibody, to yield false positive or false negative results. The selection  
5 of "blockers" therefore adds substantially to the specificity of the assays described in the present invention.

It has been found that a number of nonrelevant (i.e., nonspecific) antibodies of the same class or  
10 subclass (isotype) as those used in the assays (e.g., IgG<sub>1</sub>, IgG<sub>2a</sub>, IgM, etc.) can be used as "blockers." The concentration of the "blockers" (normally 1-100 µg/µl) is important, in order to maintain the proper sensitivity yet inhibit any unwanted interference by mutually  
15 occurring cross reactive proteins in human serum. In addition, the buffer system containing the "blockers" needs to be optimized. Preferred buffers are those based on weak organic acids, such as imidazole, HEPPS, MOPS, TES, ADA, ACES, HEPES, PIPES, TRIS, and the like, at  
20 physiological pH ranges. Somewhat less preferred buffers are inorganic buffers such as phosphate, borate or carbonate. Finally, known protease inhibitors should be added (normally at 0.01-10 µg/ml) to the buffer which contains the "blockers."

25 There are many solid phase immunoabsorbents which have been employed and which can be used in the present invention. Well known immunoabsorbents include glass, polystyrene, polypropylene, dextran, nylon and other materials, in the form of tubes, beads, and microtiter  
30 plates formed from or coated with such materials, and the like. The immobilized antibodies can be either covalently or physically bound to the solid phase immunoabsorbent, by techniques such as covalent bonding via an amide or ester linkage, or by absorption. Those  
35 skilled in the art will know many other suitable solid

- 26 -

phase immunoabsorbents and methods for immobilizing antibodies thereon, or will be able to ascertain such, using no more than routine experimentation.

For in vivo, in vitro, or in situ diagnosis, labels such as radionuclides may be bound to antibodies according to the present invention either directly or by using an intermediary functional group. An intermediary group which is often used to bind radioisotopes which exist as metallic cations to antibodies is diethylenetriaminepentaacetic acid (DTPA). Typical examples of metallic cations which are bound in this manner are:  $^{99m}\text{Tc}$ ,  $^{123}\text{I}$ ,  $^{111}\text{In}$ ,  $^{131}\text{I}$ ,  $^{97}\text{Ru}$ ,  $^{67}\text{Cu}$ ,  $^{67}\text{Ga}$  and  $^{68}\text{Ga}$ . The antibodies of the invention can also be labeled with non-radioactive isotopes for purposes of diagnosis. Elements which are particularly useful in this manner are  $^{157}\text{Gd}$ ,  $^{55}\text{Mn}$ ,  $^{162}\text{Dy}$ ,  $^{52}\text{Cr}$  and  $^{56}\text{Fe}$ .

The antigen of the invention may be isolated in substantially pure form employing antibodies according to the present invention. Thus, an embodiment of the present invention provides for substantially pure T cell receptor, B cell receptor, or Fc receptor chimera, said antigen characterized in that it is recognized by and binds to antibodies according to the present invention. In another embodiment, the present invention provides a method of isolating or purifying the receptor chimeric antigen, by forming a complex of said antigen with one or more antibodies directed against the receptor chimera.

The substantially pure T cell receptor, B cell receptor, or Fc receptor chimera antigens of the present invention may in turn be used to detect or measure antibody to the chimera in a sample, such as serum or urine. Thus, one embodiment of the present invention comprises a method of detecting the presence or amount of antibody to receptor chimera antigen in a sample, comprising contacting a sample containing an antibody to



- 27 -

the chimeric antigen with detectably labeled receptor chimera, and detecting said label. It will be appreciated that immunoreactive fractions and immunoreactive analogues of the chimera also may be used.

5 By the term "immunoreactive fraction" is intended any portion of the chimeric antigen which demonstrates an equivalent immune response to an antibody directed against the receptor chimera. By the term "immunoreactive analogue" is intended a protein which  
10 differs from the receptor chimera protein by one or more amino acids, but which demonstrates an equivalent immunoresponse to an antibody of the invention.

By "specifically recognizes and binds" is meant an antibody which recognizes and binds a chimeric receptor  
15 polypeptide but which does not substantially recognize and bind other molecules in a sample, e.g., in a biological sample, which includes the receptor polypeptide.

By "autoimmune-generated cell" is meant cells  
20 producing antibodies that react with host tissue or immune effector T cells that are autoreactive; such cells include antibodies against acetylcholine receptors (leading, e.g., to myasthenia gravis) or anti-DNA, anti-erythrocyte, and anti-platelet autoantibodies (leading,  
25 e.g., to lupus erythematosus).

By "therapeutic cell" is meant a cell which has been transformed by a chimera of the invention so that it is capable of recognizing and destroying a specific infective agent, a cell infected by a specific agent, a  
30 tumor or cancerous cell, or an autoimmune-generated cell; preferably such therapeutic cells are cells of the hematopoietic system.

By "extracellular" is meant having at least a portion of the molecule exposed at the cell surface. By  
35 "intracellular" is meant having at least a portion of the

- 28 -

molecule exposed to the therapeutic cell's cytoplasm. By "transmembrane" is meant having at least a portion of the molecule spanning the plasma membrane. An "extracellular portion", an "intracellular portion" and a "transmembrane  
5 portion", as used herein, may include flanking amino acid sequences which extend into adjoining cellular compartments.

By "oligomerize" is meant to complex with other proteins to form dimers, trimers, tetramers, or other  
10 higher order oligomers. Such oligomers may be homo-oligomers or hetero-oligomers. An "oligomerizing portion" is that region of a molecule which directs complex (i.e., oligomer) formation.

By "cytolytic" is meant to be capable of  
15 destroying a cell (e.g., a cell infected with a pathogen, a tumor or cancerous cell, or an autoimmune-generated) cell or to be capable of destroying an infective agent (e.g., a virus).

By "immunodeficiency virus" is meant a retrovirus  
20 that, in wild-type form, is capable of infecting T4 cells of a primate host and possesses a viral morphogenesis and morphology characteristic of the lentivirus subfamily. The term includes, without limitation, all variants of HIV and SIV, including HIV-1, HIV-2, SIVmac, SIVagm,  
25 SIVmnd, SIVsmm, SIVman, SIVmand, and SIVcpz.

By "MHC-independent" is meant that the cellular cytolytic response does not require the presence of an MHC class II antigen on the surface of the targeted cell.

By a "functional cytolytic signal-transducing  
30 derivative" is meant a functional derivative (as defined above) which is capable of directing at least 10%, preferably 40%, more preferably 70%, or most preferably at least 90% of the biological activity of the wild type molecule. As used herein, a "functional cytolytic  
35 signal-transducing derivative" may act by directly

- 29 -

signaling the therapeutic cell to destroy a receptor-bound agent or cell (e.g., in the case of an intracellular chimeric receptor portion) or may act indirectly by promoting oligomerization with cytolytic  
5 signal transducing proteins of the therapeutic cell (e.g., in the case of a transmembrane domain). Such derivatives may be tested for efficacy, e.g., using the in vitro assays described herein.

By a "functional HIV envelope-binding derivative"  
10 is meant a functional derivative (as defined above) which is capable of binding any HIV envelope protein. Functional derivatives may be identified using, e.g., the in vitro assays described herein.

#### THERAPEUTIC ADMINISTRATION

15 The transformed cells of the present invention may be used for the therapy of a number of diseases. Current methods of administering such transformed cells involve adoptive immunotherapy or cell-transfer therapy. These methods allow the return of the transformed immune-system  
20 cells to the bloodstream. Rosenberg, S.A., Scientific American, 62 (May 1990); Rosenberg et al., The New England Journal of Medicine, 323(9):570 (1990).

The pharmaceutical compositions of the invention may be administered to any animal which may experience  
25 the beneficial effects of the compounds of the invention. Foremost among such animals are humans, although the invention is not intended to be so limited.

#### Detailed Description

The drawings will first be described.

#### 30 Brief Description of the Drawings

FIG. 1 Characterization of CD4 chimeras. Fig. 1A presents the amino acid sequence about the site of fusion

- 30 -

between CD4 (residues 1-369) and the different receptor chains. The underlined sequence shows the position of the amino acids encoded within the BamHI site used for fusion construction. The beginning of the transmembrane domain is marked with a vertical bar. The  $\eta$  sequence is identical to the  $\zeta$  sequence at the amino terminus, but diverges at the carboxyl terminus (Jin et al., Proc. Natl. Acad. Sci. USA, 87:3319-3323 (1990)). Fig. 1B presents flow cytometric analysis of surface expression of CD4, CD4: $\zeta$ , CD4: $\gamma$  and CD4: $\eta$  in CV1 cells. Cells were infected with virus expressing CD4 chimeras or CD16<sub>PI</sub>, incubated for 9 hours at 37°C, and stained with phycoerythrin-conjugated anti-CD4 MAb Leu3A. Fig. 1C presents immunoprecipitation of labeled CD4: $\zeta$ , CD4: $\gamma$ , or native CD4 expressed in CV1 cells. Lanes were run with reducing (R) or without reducing (NR) agent. Molecular mass standards in kD are shown at left.

FIG. 2 Surface expression of CD16<sub>TM</sub> following coinfection of CD16<sub>TM</sub> alone (dense dots), or coinfecting with virus expressing CD4: $\gamma$  (dashes) or CD4: $\zeta$  (solid line). Sparse dots, cells infected with CD4: $\zeta$  alone, stained with 3G8 (Fleit et al., Proc. Natl. Acad. Sci. USA, 79:3275-3279 (1982)) (anti-CD16 MAb).

FIG. 3 Mutant CD4: $\zeta$  chimeric receptors lacking  $\zeta$  Asp-15 do not support the coexpression of CD16<sub>TM</sub>. Fig. 3A is an autoradiogram of immunoprecipitated mutant chimeras electrophoresed either with reduction (R) or without reduction (NR). Fig. 3B details surface expression of CD16<sub>TM</sub> following coinfection by viruses expressing CD16<sub>TM</sub> and the following  $\zeta$  chimeras: CD4: $\zeta$  (thick line), CD4: $\zeta$  C11G (solid line); CD4: $\zeta$  (dashed line); CD4: $\zeta$  C11G/D15G (dense dots); no coinfection (CD16<sub>TM</sub> alone, sparse dots). Cells were incubated with anti-CD16 MAb 3G8 and phycoerythrin-conjugated Fab'<sub>2</sub> goat antibodies to mouse IgG. The level of expression of the  $\zeta$  chimeras was

- 31 -

essentially identical for the different mutants analyzed, and coinfection of cells with viruses expressing CD16<sup>TM</sup> and  $\zeta$  chimeras did not appreciably alter surface expression of the chimeras (data not shown).

5           **FIG. 4** Increased intracellular free calcium ion follows crosslinking of mutant  $\zeta$  chimeras in a T cell line. Jurkat E6 cells (Weiss et al., J. Immunol., 133:123-128 (1984)) were infected with recombinant vaccinia viruses and analyzed by flow cytometry. The results shown are for the gated CD4<sup>+</sup> population, so that only cells expressing the relevant chimeric protein are analyzed. The mean ratio of violet to blue Indo-1 fluorescence reflects the intracellular free calcium concentration in the population as a whole and the percentage of responding cells reflects the fraction of cells which exceed a predetermined threshold ratio (set so that 10% of untreated cells are positive). Fig. 4A and Fig. 4B show Jurkat cells expressing CD4: $\zeta$  (solid line) or CD16: $\zeta$  (dashed line) which were exposed to anti-CD4 MAb Leu3a (phycoerythrin conjugate), followed by crosslinking with goat antibody to mouse IgG. The dotted line shows the response of uninfected cells to anti-CD3 MAb OKT3. Figs. 4C and 4D show Jurkat cells expressing CD4: $\zeta$ D15G (solid line); CD4: $\zeta$ C11G/D15G (dashes); or CD4: $\zeta$ C11G (dots) which were treated and analyzed as in Figs. 4A and 4B.

**FIG. 5** CD4: $\zeta$ , CD4: $\eta$ , and CD4: $\gamma$  receptors allow cytolytic T lymphocytes (CTL) to kill targets expressing HIV-1 gp120/41. Fig. 5A: solid circles, CTL expressing CD4: $\zeta$  incubated with HeLa cells expressing gp120/41; open circles, CTL expressing CD4: $\zeta$  incubated with uninfected HeLa cells; solid squares, uninfected CTL incubated with HeLa cells expressing gp120/41; open squares, uninfected CTL incubated with uninfected HeLa cells. Fig. 5B: solid circles, CTL expressing CD4: $\eta$  incubated with HeLa

- 32 -

cells expressing gp120/41; open circles, CTL expressing CD4: $\gamma$  incubated with HeLa cells expressing gp120/41; open squares, CTL expressing the C11G/D15G double mutant CD4: $\gamma$  chimera incubated with HeLa cells expressing gp120/41.

- 5 Fig. 5C: Flow cytometric analysis of CD4 expression by the CTL used in Fig. 5B. To correct the target to effector ratios the percent of cells expressing CD4 chimera was determined by subtracting the scaled negative (uninfected) population by histogram superposition; for  
10 comparative purposes in this figure the uninfected cells were assigned an arbitrary threshold which gives roughly the same fraction positive for the other cell populations as would histogram subtraction.

**FIG. 6 Specificity of the CD4-directed cytotoxicity.**

- 15 Fig. 6A: solid circles, CTL expressing CD4: $\gamma$  incubated with HeLa cells expressing CD16<sub>PI</sub>; open circles, CTL expressing CD4 incubated with HeLa cells expressing gp120; solid squares, CTL expressing CD16: $\gamma$  incubated with HeLa cells expressing gp120/41; open squares, CTL  
20 expressing CD16<sub>PI</sub> incubated with HeLa cells expressing gp120/41. Fig. 6B: solid circles, CTL expressing CD4: $\gamma$  incubated with Raji (MHC class II<sup>+</sup>) cells; open circles, uninfected CTL cells incubated with RJ2.2.5 (MHC class II<sup>-</sup> Raji mutant) cells; solid squares, uninfected CTL  
25 incubated with Raji (MHC class II<sup>+</sup>) cells; open squares, CTL expressing CD4: $\gamma$  incubated with RJ2.2.5 (MHC class II<sup>-</sup>) cells. The ordinate scale is expanded.

- FIG. 7 Characterization of the CD16: $\gamma$  chimeric receptor.** Fig. 7A is a schematic diagram of the CD16: $\gamma$   
30 fusion protein. The extracellular portion of the phosphatidylinositol-linked form of monomeric CD16 was joined to dimeric  $\gamma$  just external to the transmembrane domain. The protein sequence at the fusion junction is shown at the bottom. Fig. 7B shows a flow cytometric  
35 analysis of calcium mobilization following crosslinking

- 33 -

of the CD16: $\zeta$  chimera in either a TCR positive or TCR positive or TCR negative cell line. The mean ratio of violet to blue fluorescence (a measure of relative calcium ion concentration) among cell populations treated with antibodies at time 0 is shown. Solid squares, the response of Jurkat cells to anti-CD3 MAb OKT3; solid triangles, the response of CD16: $\zeta$  to anti-CD16 MAb 3G8 crosslinking in the REX33A TCR<sup>-</sup> mutant; open squares, the response to CD16: $\zeta$  crosslinking in the Jurkat TCR<sup>-</sup> mutant line JRT3.T3.5; open triangles, the response to CD16: $\zeta$  crosslinking in Jurkat cells; crosses, the response to nonchimeric CD16 in Jurkat cells; and dots, the response to nonchimeric CD16 in the REX33A TCR<sup>-</sup> cell line.

**FIG. 8 Deletion analysis of cytolytic potential.**

Fig. 8A shows the locations of the  $\zeta$  deletion endpoints. Here as elsewhere mutations in  $\zeta$  are represented by the original residue-location-mutant residue convention, so that D66\*, for example, denotes replacement of Asp-66 by a termination codon. Fig. 8B shows cytolysis assay results of undeleted CD16: $\zeta$  and salient  $\zeta$  deletions. Hybridoma cells expressing surface antibody to CD16 were loaded with <sup>51</sup>Cr and incubated with increasing numbers of human cytolytic lymphocytes (CTL) infected with vaccinia recombinants expressing CD16: $\zeta$  chimeras. The percent of <sup>51</sup>Cr released is plotted as a function of the effector (CTL) to target (hybridoma) cell ratio (e/t). Solid circles, cytolysis mediated by cells expressing CD16: $\zeta$  (mfi 18.7); solid squares, cytolysis mediated by cells expressing CD16: $\zeta$  Asp66\* (mfi 940.2); open squares, cytolysis mediated by cells expressing CD16: $\zeta$  Glu60\* (mfi 16.0); open circles, cytolysis mediated by cells expressing CD16: $\zeta$  Tyr51\* (mfi 17.4); solid triangles, cytolysis mediated by cells expressing CD16: $\zeta$  Phe34\* (mfi 17.8); and open triangles, cytolysis mediated by cells expressing nonchimeric CD16 (mfi 591). Although in this

- 34 -

experiment the expression of CD16:ζAsp66\* was not matched to that of the other fusion proteins, cytolysis by cells expressing CD16:ζ at equivalent levels in the same experiment gave results essentially identical to those shown by cells expressing CD16:ζAsp66\* (not shown).

**FIG. 9** Elimination of the potential for transmembrane interactions reveals a short ζ segment capable of mediating cytolysis. Fig. 9A is a schematic diagram of the monomeric bipartite and tripartite chimeras. At the top is the CD16:ζ construct truncated at residue 65 and lacking transmembrane Cys and Asp residues. Below are the CD16:CD5:ζ and CD16:CD7:ζ constructs and related controls. The peptide sequences of the intracellular domains are shown below. Fig. 9B shows the cytolytic activity of monomeric chimera deletion mutants. The cytolytic activity of cells expressing CD16:ζ (solid circles; mfi 495) was compared to that of cells expressing CD16:ζAsp66\* (solid squares; mfi 527) or the mutants CD16:ζCys11Gly/Asp15Gly/Asp66\*, (open squares; mfi 338) and CD16:ζCys11Gly/Asp15Gly/Glu60\* (filled triangles; mfi 259). Fig. 9C shows the cytolytic activity mediated by tripartite fusion proteins. Solid triangles, CD16:ζAsp66\*; open squares, CD16:5:ζ(48-65); solid squares CD16:7:ζ(48-65); open triangles, CD16:7:ζ(48-59); open circles, CD16:5; solid circles, CD16:7. Fig. 9D shows calcium mobilization by mutant and tripartite chimeras in the TCR negative Jurkat JRT3.T3.5 mutant cell line. Open circles, response of cells expressing dimeric CD16:ζAsp66\*; solid squares, response of cells expressing CD16:ζCys11Gly/Asp15Gly/Asp66\*; open squares, response of cells expressing CD16:ζCys11Gly/Asp15Gly/Glu60\*; solid triangles, response of cells expressing CD16:7:ζ(48-65); and open triangles, response of cells expressing CD16:ζ(48-59).



- 35 -

FIG. 10 Contribution of individual amino acids to the activity of the 18 residue cytolytic signal-transducing motif. Figs. 10A and 10B show cytolytic activity and Fig. 10C shows calcium ion mobilization mediated by chimeras bearing point mutations near the carboxyl terminal tyrosine (Y62). Figs. 10A and 10B represent data collected on cells expressing low and high amounts, respectively, of the CD16: $\zeta$  fusion proteins. Identical symbols are used for the calcium mobilization and cytolysis assays, and are shown in one letter code at right. Solid circles, cells expressing CD16: $\zeta$  (mfi in A, 21; B, 376); solid squares, cells expressing CD16:7: $\zeta$ (48-65) (mfi A, 31; B, 82); open squares, CD16:7: $\zeta$ (48-65)Glu60Gln (mfi A, 33; B, 92), crosses, CD16:7: $\zeta$ (48-65)Asp63Asn (mfi A, 30; B, 74); solid triangles, CD16:7: $\zeta$ (48-65)Tyr62Phe (mfi A, 24; B, 88); open circles, CD16:7: $\zeta$ (48-65)Glu61Gln (mfi A, 20; B, 62); and open triangles, CD16:7: $\zeta$ (48-65)Tyr62Ser (mfi B, 64). Figs. 10D and 10E show cytolytic activity and Fig. 10F shows calcium ion mobilization by chimeras bearing point mutations near the amino terminal tyrosine (Y51). Identical symbols are used for the calcium mobilization and cytolysis assays and are shown at right. Solid circles, cells expressing CD16: $\zeta$  (mfi in D, 21.2; in E, 672); solid squares, cells expressing CD16:7: $\zeta$ (48-65) (mfi D, 31.3; E, 179); solid triangles, CD16:7: $\zeta$ (48-65)Asn48Ser (mfi D, 22.4; E, 209); open squares, CD16:7: $\zeta$ (48-65)Leu50Ser (mfi D, 25.0; E, 142); and open triangles, CD16:7: $\zeta$ (48-65)Tyr51Phe (mfi D, 32.3; E, 294).

FIG. 11 Alignment of internal repeats of  $\zeta$  and comparison of their ability to support cytolysis. Fig. 11A is a schematic diagram of chimeras formed by dividing the  $\zeta$  intracellular domain into thirds and appending them to the transmembrane domain of a CD16:7 chimera. The sequences of the intracellular domains are shown below,

- 36 -

with shared residues boxed, and related residues denoted by asterisks. Fig. 11B shows the cytolytic potency of the three  $\zeta$  subdomains. Solid circles, cells expressing CD16: $\zeta$  (mfi 476); solid squares, CD16:7: $\zeta$ (33-65) (mfi 68); open squares, CD16:7: $\zeta$ (71-104) (mfi 114); and solid triangles, CD16:7: $\zeta$ (104-138) (mfi 104).

FIG. 12 is a schematic diagram of the CD16:FcR $\gamma$ II chimeras.

FIG. 13 Calcium mobilization following crosslinking of CD4:FcR $\gamma$ II and CD16:FcR $\gamma$ II chimeras. Fig. 13A shows the ratio of violet to blue fluorescence emitted by cells loaded with the calcium sensitive fluorophore Indo-1 shown as a function of time following crosslinking of the CD16 extracellular domain with antibodies. Fig. 13B shows a similar analysis of the increase in ratio of violet to blue fluorescence of cells bearing CD4:FcR $\gamma$ II chimeras, following crosslinking with antibodies.

FIG. 14 Cytolysis assay of CD4:FcR $\gamma$ II and CD16:FcR $\gamma$ II chimeras. Fig. 14A shows the percent of  $^{51}\text{Cr}$  released from anti-CD16 hybridoma (target) cells when the cells are exposed to increasing numbers of cytotoxic T lymphocytes expressing CD16:FcR $\gamma$ II chimeras (effector cells). Fig. 14B shows a similar analysis of cytotoxicity mediated by CD4:FcR $\gamma$ II chimeras against target cells expressing HIV envelope glycoproteins.

FIG. 15 Identification of residues in the FcR $\gamma$ II A tail which are important for cytolysis. Fig. 15A is a schematic diagram of the deletion constructs. Figs. 15B and 15C shows calcium mobilization and cytolysis by carboxyl-terminal deletion variants of CD16:FcR $\gamma$ II A. Figs. 15D and 15E show calcium mobilization and cytolysis by tripartite chimeras bearing progressively less of the amino terminus of the intracellular tail of CD16:FcR $\gamma$ II A.

- 37 -

FIG. 16 ( SEQ ID NO: 24) shows the amino acid sequence of the CD3 delta receptor protein; the boxed sequence represents a preferred cytolytic signal transducing portion.

5        FIG. 17 (SEQ ID NO: 25) shows the amino acid sequence of the T3 gamma receptor protein; the boxed sequence represents a preferred cytolytic signal transducing portion.

10       FIG. 18 (SEQ ID NO: 26) shows the amino acid sequence of the mb1 receptor protein; the boxed sequence represents a preferred cytolytic signal transducing portion.

15       FIG. 19 (SEQ ID NO: 27) shows the amino acid sequence of the B29 receptor protein; the boxed sequence represents a preferred cytolytic signal transducing portion.

#### EXAMPLE I

##### Construction of Human IgG1:Receptor Chimeras

Human IgG1 heavy chain sequences were prepared by joining sequences in the C<sub>H</sub>3 domain to a cDNA fragment derived from the 3' end of the transmembrane form of the antibody mRNA. The 3' end fragment was obtained by polymerase chain reaction using a tonsil cDNA library as substrate, and oligonucleotides having the sequences:

25        CGC GGG GTG ACC GTG CCC TCC AGC AGC TTG GGC (SEQ ID NO: 7) and

CGC GGG GAT CCG TCG TCC AGA GCC CGT CCA GCT CCC CGT CCT GGG CCT CA (SEQ ID NO: 8), corresponding to the 5' and 3' ends of the desired DNA fragments respectively. The 5' oligo is complementary to a site in the C<sub>H</sub>1 domain of human IgG1, and the 3' oligo is complementary to a site just 5' of the sequences encoding the membrane spanning domain. The PCR product was digested with BstXI and BamHI and ligated between

30

- 38 -

BstXI and BamHI sites of a semisynthetic IgG1 antibody gene bearing variable and constant regions. Following the insertion of the BstXI to BamHI fragment, the amplified portions of the construct were replaced up to the SmaI site in C<sub>H</sub>3 by restriction fragment interchange, so that only the portion between the SmaI site and the 3' oligo was derived from the PCR reaction.

To create a human IgG1: $\zeta$  chimeric receptor, the heavy chain gene ending in a BamHI site was joined to the BamHI site of the  $\zeta$  chimera described below, so that the antibody sequences formed the extracellular portion. Flow cytometry of COS cells transfected with a plasmid encoding the chimera showed high level expression of antibody determinants when an expression plasmid encoding a light chain cDNA was cotransfected, and modest expression of antibody determinants when the light chain expression plasmid was absent.

Similar chimeras including human IgG1 fused to  $\eta$  or  $\gamma$  (see below), or any signal-transducing portion of a T cell receptor or Fc receptor protein may be constructed generally as described above using standard techniques of molecular biology.

To create a single transcription unit which would allow both heavy and light chains to be expressed from a single promoter, a plasmid encoding a bicistronic mRNA was created from heavy and light chain coding sequences, and the 5' untranslated portion of the mRNA encoding the 78kD glucose regulated protein, otherwise known as grp78, or BiP. grp78 sequences were obtained by PCR of human genomic DNA using primers having the sequences:

CGC GGG CGG CCG CGA CGC CGG CCA AGA CAG CAC (SEQ ID NO: 9) and

CGC GTT GAC GAG CAG CCA GTT GGG CAG CAG CAG (SEQ ID NO: 10)

- 39 -

at the 5' and 3' ends respectively. Polymerase chain reactions with these oligos were performed in the presence of 10% dimethyl sulfoxide. The fragment obtained by PCR was digested with NotI and HincII and  
5 inserted between NotI and HpaI sites downstream from human IgG1 coding sequences. Sequences encoding a human IgG kappa light chain cDNA were then inserted downstream from the grp78 leader, using the HincII site and another site in the vector. The expression plasmid resulting  
10 from these manipulations consisted of the semisynthetic heavy chain gene, followed by the grp78 leader sequences, followed by the kappa light chain cDNA sequences, followed by polyadenylation signals derived from an SV40 DNA fragment. Transfection of COS cells with the  
15 expression plasmid gave markedly improved expression of heavy chain determinants, compared to transfection of plasmid encoding heavy chain determinants alone.

To create a bicistronic gene comprising a heavy chain/receptor chimera and a light chain, the upstream  
20 heavy chain sequences can be replaced by any chimeric heavy chain/ receptor gene described herein.

## EXAMPLE II

### Construction of CD4 Receptor Chimeras

Human  $\zeta$  (Weissman et al., Proc. Natl. Acad. Sci. USA, 85:9709-9713 (1988b)) and  $\gamma$  (Küster et al., J. Biol. Chem., 265:6448-6452 (1990)) cDNAs were isolated by  
25 polymerase chain reaction from libraries prepared from the HPB-ALL tumor cell line (Aruffo et al., Proc. Natl. Acad. Sci. USA, 84:8573-8577 (1987b)) and from human  
30 natural killer cells, while  $\eta$  cDNA (Jin et al., Proc. Natl. Acad. Sci. USA, 87:3319-3323 (1990)) was isolated from a murine thymocyte library.  $\zeta$ ,  $\eta$  and  $\gamma$  cDNAs were joined to the extracellular domain of an engineered form of CD4 possessing a BamHI site just upstream of the

- 40 -

membrane spanning domain (Aruffo et al., Proc. Natl. Acad. Sci. USA, 84:8573-8577 (1987b); Zettlmeissl et al., DNA Cell Biol., 9:347-353 (1990)) which was joined to the BamHI site naturally present in the  $\zeta$  and  $\eta$  cDNAs at a similar location a few residues upstream of the membrane spanning domain (SEQ ID NOS: 1, 3, 4 and 6). To form the fusion protein with  $\gamma$  a BamHI site was engineered into the sequence at the same approximate location (Fig. 1; SEQ ID NO: 2 and 5). The gene fusions were introduced into a vaccinia virus expression plasmid bearing the E. coli gpt gene as a selectable marker (M. Amiot and B.S., unpublished), and inserted into the genome of the vaccinia WR strain by homologous recombination and selection for growth in mycophenolic acid (Falkner et al., J. Virol., 62:1849-1854 (1988); Boyle et al., Gene, 65:123-128 (1988)). Flow cytometric analysis showed that the vaccinia recombinants direct the abundant production of CD4: $\zeta$  and CD4: $\gamma$  fusion proteins at the cell surface, whereas the expression of CD4: $\eta$  is substantially weaker (Fig. 1). The latter finding is consistent with a recent report that transfection of an  $\eta$  cDNA expression plasmid into a murine hybridoma cell line gave substantially less expression than transfection of a comparable  $\zeta$  expression plasmid (Clayton et al., J. Exp. Med., 172:1243-1253 (1990)). Immunoprecipitation of cells infected with the vaccinia recombinants revealed that the fusion proteins form covalent dimers, unlike the naturally occurring CD4 antigen (Fig. 1). The molecular masses of the monomeric CD4: $\zeta$  and CD4: $\gamma$  fusion proteins and native CD4 were found to be 63, 55 and 53 kD respectively. The larger masses of the fusion proteins are approximately consistent with the greater length of the intracellular portion, which exceeds that of native CD4 by 75 (CD4: $\zeta$ ) or 5 (CD4: $\gamma$ ) residues.

- 41 -

## EXAMPLE III

## CD4 Chim ras Can Associat With Other Receptor Chains

Cell surface expression of the macrophage/natural killer cell form of human Fc $\gamma$ RIII (CD16<sub>TM</sub>) on  
5 transfectants is facilitated by cotransfection with murine (Kurosaki et al., Nature, 342:805-807 (1989)) or human (Hibbs et al., Science, 246:1608-1611 (1989))  $\gamma$ , as well as by human  $\zeta$  (Lanier et al., Nature, 342:803-805 (1989)).

10 Consistent with these reports, expression of the chimeras also allowed surface expression of CD16<sub>TM</sub> when delivered to the target cell either by cotransfection or by coinfection with recombinant vaccinia viruses (Fig. 2). The promotion of (CD16<sub>TM</sub>) surface expression by  $\zeta$  was  
15 more pronounced than promotion by  $\gamma$  (Fig. 2) in the cell lines examined, whereas native CD4 (data not shown) did not enhance CD16<sub>TM</sub> surface expression.

## EXAMPLE IV

Asp  $\zeta$  Mutants Do Not Coassociate with Fc Receptor

20 To create chimeras which would not associate with existing antigen or Fc receptors, mutant  $\zeta$  fusion proteins which lacked either the intramembranous Asp or intramembranous Cys residue or both were prepared. Flow cytometry showed that the intensity of cell surface  
25 expression by the different mutant chimeras was not appreciably different from the unmutated precursor (data not shown) and immunoprecipitation experiments showed that total expression by the chimeras was similar (Fig. 3). As expected, the mutant chimeras lacking the  
30 transmembrane cysteine residue were found not to form disulfide linked dimers (Fig. 3). The two mutant chimeras lacking Asp were incapable of supporting the surface expression of CD16<sup>m</sup>, whereas the monomeric chimeras lacking Cys but bearing Asp allowed CD16<sub>TM</sub> to be

- 42 -

coexpressed, but at lower efficiency than the parental dimer (Fig. 3).

#### EXAMPLE V

##### Mutant Receptors Retain the Ability to Initiate a Calcium Response

To determine whether crosslinking of the fusion proteins would allow the accumulation of free intracellular calcium in a manner similar to that known to occur with the T cell antigen receptor, cells of the human T cell leukemia line, Jurkat E6 (ATCC Accession Number TIB 152, American Type Culture Collection, Rockville, MD), were infected with the vaccinia recombinants and the relative cytoplasmic calcium concentration following crosslinking of the extracellular domain with antibodies was measured. Flow cytometric measurements were performed with cells loaded with the calcium sensitive dye Indo-1 (Grynkiewicz et al., J. Biol. Chem., 260:3340-3450 (1985); Rabinovitch et al., J. Immunol., 137:952-961 (1986)). Figure 4 shows the results of calcium flux experiments with cells infected with CD4: $\zeta$  and the Asp<sup>-</sup> and Cys<sup>-</sup> mutants of  $\zeta$ . Crosslinking of the chimeras, reproducibly increased intracellular calcium. CD4: $\eta$  and CD4: $\gamma$  similarly allowed accumulation intracellular calcium in infected cells (data not shown). Jurkat cells express low levels of CD4 on the cell surface, however, crosslinking of the native CD4 in the presence or absence of CD16: $\zeta$  (C.R. and B.S. unpublished) (Fig. 4 and data not shown) does not alter intracellular calcium levels.

#### EXAMPLE VI

##### CD4: $\zeta$ , $\eta$ , and $\gamma$ Chimeras Mediate Cytolysis of Targets Expressing HIV gp120/41

To determine whether the chimeric receptors would trigger cytolytic effector programs, a model



- 43 -

target:effector system based on CD4 recognition of the HIV envelope gp120/gp41 complex was created. HeLa cells were infected with recombinant vaccinia viruses expressing gp120/gp41 (Chakrabarti et al., Nature, 5 320:535-537 (1986); Earl et al., J. Virol., 64:2448-2451 (1990)) and labeled with  $^{51}\text{Cr}$ . The labeled cells were incubated with cells from a human allospecific ( $\text{CD8}^+$ ,  $\text{CD4}^-$ ) cytotoxic T lymphocyte line which had been infected with vaccinia recombinants expressing the  $\text{CD4}:\zeta$ ,  $\text{CD4}:\eta$ , 10 or  $\text{CD4}:\gamma$  chimeras, or the  $\text{CD4}:\zeta\text{Cys11Gly:Asp15Gly}$  double mutant chimera. Fig. 5 shows that HeLa cells expressing gp120/41 were specifically lysed by cytotoxic T lymphocytes (CTL) expressing CD4 chimeras. Uninfected HeLa cells were not targeted by CTL armed with  $\text{CD4}:\zeta$  15 chimeras, and HeLa cells expressing gp120/41 were not recognized by uninfected CTL. To compare the efficacy of the various chimeras, the effector to target ratios were corrected for the fraction of CTL expressing CD4 chimeras, and for the fraction of HeLa cells expressing 20 gp120/41, as measured by flow cytometry. Fig. 5C shows a cytometric analysis of CD4 expression by the CTL used in the cytolysis experiment shown in Figs. 4A and 4B. Although the mean density of surface  $\text{CD4}:\zeta$  greatly exceeded the mean density of  $\text{CD4}:\eta$ , the cytolytic 25 efficiencies of cells expressing either form were similar. Correcting for the fraction of targets expressing gp120, the efficiency of cytolysis mediated by  $\text{CD4}:\zeta$  and  $\text{CD4}:\eta$  proteins are comparable to the best efficiencies reported for specific T cell receptor 30 target:effector pairs (the mean effector to target ratio for 50% release by T cells expressing  $\text{CD4}:\zeta$  was  $1.9 \pm 0.99$ ,  $n=10$ ). The  $\text{CD4}:\gamma$  fusion was less active, as was the  $\text{CD4}:\zeta$  fusion lacking the transmembrane Asp and Cys residues. However in both cases significant cytolysis 35 was observed (Fig. 5).

- 44 -

To control for the possibility that vaccinia infection might promote artefactual recognition by CTL, similar cytolysis experiments were performed with target cells infected with vaccinia recombinants expressing the phosphatidylinositol linked form of CD16 (CD16<sub>PI</sub>) and labeled with <sup>51</sup>Cr, and with CTL infected with control recombinants expressing either CD16<sub>PI</sub> or CD16:ζ. Fig. 6A shows that T cells expressing non-CD4 chimeras do not recognize native HeLa cells or HeLa cells expressing gp120/41, and similarly that T cells expressing CD4 chimeras do not recognize HeLa cells expressing other vaccinia-encoded surface proteins. In addition, CTLs expressing non-chimeric CD4 do not significantly lyse HeLa cells expressing gp120/41 (Fig. 6A).

15

#### EXAMPLE VII

##### MHC Class II-Bearing Cells Are Not Targeted by the Chimeras

CD4 is thought to interact with a nonpolymorphic sequence expressed by MHC class II antigen (Gay et al., Nature, 328:626-629 (1987); Sleckman et al., Nature, 328:351-353 (1987)). Although a specific interaction between CD4 and class II antigen has never been documented with purified proteins, under certain conditions adhesion between cells expressing CD4 and cells expressing class II molecules can be demonstrated (Doyle et al., Nature, 330:256-259 (1987); Clayton et al., J. Exp. Med., 172:1243-1253 (1990); Lamarre et al., Science, 245:743-746 (1989)). Next examined was whether killing could be detected against cells bearing class II. Fig. 6B shows that there is no specific cytolysis directed by CD4:ζ against the Raji B cell line, which expresses abundant class II antigen. Although a modest (~5%) cytolysis is observed, a class II-negative mutant of Raji, RJ2.2.5, (Accolla, R.S., J. Exp. Med., 157:1053-

- 45 -

1058 (1983)) shows a similar susceptibility, as do Raji cells incubated with uninfected T cells.

#### EXAMPLE VIII

##### Sequence Requirements for Induction of Cytolysis by the T 5 Cell Antigen/Fc Receptor Zeta Chain

Although chimeras between CD4 and  $\zeta$  can arm cytotoxic T lymphocytes (CTL) to kill target cells expressing HIV gp120, an alternative to CD4 was sought in order to unambiguously compare the properties of zeta  
10 chimeras introduced into human T cell lines. Such lines can express CD4, making it difficult to specifically define the relationship between the type or degree of calcium mobilization and the cytotoxic potential of the different chimeras. To circumvent this, chimeras were  
15 created between  $\zeta$  and CD16 in which the extracellular domain of CD16 is attached to the transmembrane and intracellular sequences of  $\zeta$  (Fig. 7A). The gene fusions were introduced into a vaccinia virus expression plasmid bearing the E. coli gpt gene as a selectable marker and  
20 inserted into the genome of the vaccinia WR strain by homologous recombination and selection for growth in mycophenolic acid (Falkner and Moss, J. Virol. 62:1849 (1988); Boyle and Coupar, Gene 65:123 (1988)).

T cell lines were infected with the vaccinia  
25 recombinants and the relative cytoplasmic free calcium ion concentration was measured following crosslinking of the extracellular domains with antibodies. Both spectrofluorimetric (bulk population) and flow cytometric (single cell) measurements were performed, with cells  
30 loaded with the dye Indo-1 (Grynkiewicz et al., J. Biol. Chem. 260:3440 (1985); Rabinovitch et al., J. Immunol. 137:952 (1986)). Figure 7B shows an analysis of data collected from cells of the Jurkat human T cell leukemia line infected with vaccinia recombinants expressing

- 46 -

CD16:ζ fusion protein. Crosslinking of the chimeras reproducibly increased intracellular calcium, while similar treatment of cells expressing nonchimeric CD16 had little or no effect. When the chimera was expressed  
5 in mutant cell lines lacking antigen receptor, either REX33A (Breitmeyer et al. J. Immunol. 138:726 (1987); Sancho et al. J. Biol. Chem. 264:20760 (1989)), or Jurkat mutant JRT3.T3.5 (Weiss et al., J. Immunol. 135:123 (1984)); or a strong response to CD16 antibody  
10 crosslinking was seen. Similar data have been collected on the REX20A (Breitmeyer et al., supra, 1987; Blumberg et al., J. Biol. Chem. 265:14036 (1990)) mutant cell line, and a CD3/Ti negative mutant of the Jurkat cell line established in this laboratory (data not shown).  
15 Infection with recombinants expressing CD16:ζ did not restore the response to anti-CD3 antibody, showing that the fusion protein did not act by rescuing intracellular CD3 complex chains (data not shown).

To evaluate the ability of the chimeras to  
20 redirect cell-mediated immunity, CTLs were infected with vaccinia recombinants expressing CD16 chimeras and used to specifically lyse hybridoma cells expressing membrane-bound anti-CD16 antibodies (see below). This assay is an extension of a hybridoma cytotoxicity assay originally  
25 developed to analyze effector mechanisms of cells bearing Fc receptors (Graziano and Fanger, J. Immunol. 138:945, 1987; Graziano and Fanger, J. Immunol. 139:35-36, 1987; Shen et al., Mol. Immunol. 26:959, 1989; Fanger et al., Immunol. Today 10: 92, 1989). Fig. 8B shows that  
30 expression of CD16:ζ in cytotoxic T lymphocytes allows the armed CTL to kill 3G8 (anti-CD16; Fleit et al., Proc. Natl. Acad. Sci. USA 79:3275, 1982) hybridoma cells, whereas CTL expressing the phosphatidylinositol-linked form of CD16 are inactive. CTL armed with CD16:ζ also do

- 47 -

not kill hybridoma cells expressing an irrelevant antibody (data not shown).

To identify the minimal  $\zeta$  sequences necessary for cytolysis, a series of deletion mutants were prepared in which successively more of the  $\zeta$  intracellular domain was removed from the carboxyl terminus (Fig. 8A). Most of the intracellular domain of zeta could be removed with little consequence for cytolytic potential; the full length chimera CD16: $\zeta$  was essentially equal in efficacy to the chimera deleted to residue 65, CD16: $\zeta$ Asp66\* (Fig. 8B). A substantial decrease in cytotoxicity was observed on deletion to  $\zeta$  residue 59 (chimera CD16: $\zeta$ Glu60\*), and further deletion to residue 50 resulted in slightly less activity. However complete loss of activity was not observed even when the intracellular domain was reduced to a three residue transmembrane anchor (Fig. 8B).

Because  $\zeta$  is a disulfide linked dimer, one explanation for the retention of cytolytic activity was that endogenous  $\zeta$  was forming heterodimers with the chimeric  $\zeta$  deletion, thereby reconstituting activity. To test this idea,  $\zeta$  residues 11 and 15 were changed from Asp and Cys respectively to Gly (Cys11Gly/Asp15Gly), and immunoprecipitations were carried out as follows. Approximately  $2 \times 10^6$  CV1 cells were infected for one hour in serum free DME medium with recombinant vaccinia at a multiplicity of infection (moi) of at least ten. Six to eight hours post-infection, the cells were detached from the plates with PBS/1mM EDTA and surface labeled with 0.2 mCi  $^{125}\text{I}$  per  $2 \times 10^6$  cells using lactoperoxidase and  $\text{H}_2\text{O}_2$  by the method of Clark and Einfeld (*Leukocyte Typing II*, pp. 155-167, Springer-Verlag, NY, 1986). The labeled cells were collected by centrifugation and lysed in 1% NP-40, 0.1% SDS, 0.15M NaCl, 0.05M Tris, pH 8.0, 5mM  $\text{MgCl}_2$ , 5mM KCl, 0.2M iodoacetamide and 1mM PMSF. Nuclei were removed by

- 48 -

centrifugation, and CD16 proteins were immunoprecipitated with antibody 3G8 (Fleit et al., supra, 1982; Medarex) and anti-mouse IgG agarose (Cappel, Durham, NC). Samples were electrophoresed through an 8% polyacrylamide/SDS gel under non-reducing conditions or through a 10% gel under reducing conditions. These immunoprecipitations confirmed that the CD16:ζCys11Gly/Asp15Gly chimera did not associate in disulfide-linked dimer structures.

The cytolytic activity of the mutant receptors was also tested. The mutated chimera deleted to residue 65 (CD16:ζCys11Gly/Asp15Gly/Asp66\*) was, depending on the conditions of assay, two to eight fold less active in the cytotoxicity assay than the comparable unmutated chimera (CD16:ζAsp66\*), which was usually within a factor of two of, or indistinguishable in activity from, CD16:ζ (Fig. 9B). The reduction in activity of the mutant chimeras is comparable to the reduction seen with CD4 chimeras of similar structure (see above) and is most likely attributable to the lower efficiency of ζ monomers compared to dimers. In contrast, the Asp<sup>-</sup>, Cys<sup>-</sup> mutated chimera deleted to residue 59 had no cytolytic activity (Fig. 9B), supporting the hypothesis that association with other chains mediated by the transmembrane Cys and/or Asp residues was responsible for the weak persistence of cytolytic activity in deletions more amino terminal than residue 65.

Flow cytometric studies showed that the deletion mutants lacking transmembrane Asp and Cys residues could still promote an increase in free intracellular calcium ion in response to antibody crosslinking in a TCR<sup>-</sup> mutant Jurkat cell line (Fig. 9D). Similar results were obtained for chimeras expressed in the parental Jurkat line (not shown). In the case of CD16:ζCys11Gly/Asp15Gly/Glu60\*, these data demonstrate that the ability to mediate calcium responsiveness can be

- 49 -

mutationally separated from the ability to support cytolysis.

To definitively eliminate the possible contribution of  $\zeta$  transmembrane residues, the transmembrane and first 17 cytoplasmic residues of  $\zeta$  were replaced by sequences encoding the membrane spanning and first 14 or first 17 cytoplasmic residues of the CD5 or CD7 antigens, respectively (Fig. 9A). The resulting tripartite fusion proteins CD16:5: $\zeta$ (48-65) and CD16:7: $\zeta$ (48-65) did not form disulfide-linked dimers as do the simpler CD16: $\zeta$  chimeras, because they lacked the cysteine residue in the  $\zeta$  transmembrane domain. Both tripartite chimeras were able to mobilize calcium in Jurkat and TCR negative cell lines (Fig. 9D) and to mount a cytolytic response in CTL (Fig. 9C and data not shown). However truncation of the  $\zeta$  portion to residue 59 in chimera CD16:7: $\zeta$ (48-59) abrogates the ability of tripartite fusion to direct calcium responsiveness in TCR positive or negative Jurkat cells or cytolysis in mature CTL (Fig. 9C and 9D and data not shown).

To examine the contributions of individual residues within the 18-residue motif, we prepared a number of mutant variants by site-directed mutagenesis, and evaluated their ability to mediate receptor-directed killing under conditions of low (Figs. 10A and 10D) or high (Figs. 10B and 10E) expression of chimeric receptor. Fig. 10 shows that while a number of relatively conservative substitutions (i.e., replacing acidic residues with their cognate amides, or tyrosine with phenylalanine) which spanned residues 59 to 63 yielded moderate compromise of cytolytic efficacy, in general the variants retained the ability to mobilize calcium. However collectively these residues comprise an important submotif inasmuch as their deletion eliminates cytolytic activity. Conversion of Tyr 62 to either Phe or Ser

- 50 -

eliminated both the cytotoxic and calcium responses. At the amino terminus of the 18 residue segment, replacement of Tyr 51 with Phe abolished both calcium mobilization and cytolytic activity, while substitution of Leu with Ser at position 50 eliminated the calcium response while only partially impairing cytolysis. Without being bound to a particular hypothesis, it is suspected that the inability of the Leu50Ser mutant to mobilize calcium in short term flow cytometric assays does not fully reflect its ability to mediate a substantial increase in free intracellular calcium ion over the longer time span of the cytolysis assay. However, calcium-insensitive cytolytic activity has been reported for some cytolytic T cell lines, and the possibility that a similar phenomenon underlies the results described herein has not been ruled out. Replacement of Asn48 with Ser partially impaired cytotoxicity in some experiments while having little effect in others.

To investigate the potential role of redundant sequence elements, the intracellular domain of  $\zeta$  was divided into three segments, spanning residues 33 to 65, 71 to 104, and 104 to 138. Each of these segments was attached to a CD16:CD7 chimera by means of a MluI site introduced just distal to the basic membrane anchoring sequences of the intracellular domain of CD7 (see below; Fig. 11A). Comparison of the cytolytic efficacy of the three elements showed they were essentially equipotent (Fig. 11B). Sequence comparison (Fig. 11A) shows that the second motif bears eleven residues between tyrosines, whereas the first and third motifs bear ten.

Although a precise accounting of the process of T cell activation has not been made, it is clear that aggregation of the antigen receptor, or of receptor chimeras which bear  $\zeta$  intracellular sequences, triggers calcium mobilization, cytokine and granule release, and



- 51 -

the appearance of cell surface markers of activation. The active site of  $\zeta$ , a short linear peptide sequence probably too small to have inherent enzymatic activity, likely interacts with one or at most a few proteins to mediate cellular activation. It is also clear that mobilization of free calcium is not by itself sufficient for cellular activation, as the ability to mediate cytolysis can be mutationally separated from the ability to mediate calcium accumulation.

As shown herein, addition of 18 residues from the intracellular domain of  $\zeta$  to the transmembrane and intracellular domain of two unrelated proteins allows the resulting chimeras to redirect cytolytic activity against target cells which bind to the extracellular portion of the fusion proteins. Although chimeras bearing the 18 residue motif are approximately eight-fold less active than chimeras based on full length  $\zeta$ , the reduced activity can be attributed to the loss of transmembrane interactions which normally allow wild-type  $\zeta$  to form disulfide linked dimers. That is,  $\zeta$  deletion constructs which have the same carboxyl terminus as the motif and lack transmembrane Cys and Asp residues typically show slightly less activity than chimeras bearing only the 18 residue motif.

The cytolytic competency element on which we have focused has two tyrosines and no serines or threonines, restricting the possible contributions of phosphorylation to activity. Mutation of either tyrosine destroys activity, however, and although preliminary experiments do not point to a substantial tyrosine phosphorylation following crosslinking of chimeric surface antigens bearing the 18 residue motif, the possible participation of such phosphorylation at a low level cannot be excluded. In addition to the effects noted at the two tyrosine residues, a number of amino acid replacements at

- 52 -

the amino and carboxyl terminus of the motif weaken activity under conditions of low receptor density.

Sequences similar to the  $\zeta$  active motif can be found in the cytoplasmic domains of several other  
5 transmembrane proteins, including the CD3  $\delta$  and  $\gamma$  molecules, the surface IgM associated proteins mb1 and B29, and the  $\beta$  and  $\gamma$  chains of the high affinity IgE receptor, Fc $\epsilon$ RI (Reth, Nature 338:383, 1989). Although the function of these sequences is uncertain, if  
10 efficiently expressed, each may be capable of autonomous T cell activation, and such activity may explain the residual TCR responsiveness seen in a zeta-negative mutant cell line (Sussman et al., Cell 52:85, 1988).

$\zeta$  itself bears three such sequences, approximately  
15 equally spaced, and a rough trisection of the intracellular domain shows that each is capable of initiating a cytolytic response.  $\eta$ , a splice isoform of  $\zeta$  (Jin et al., supra, 1990; Clayton et al., Proc. Natl. Acad. Sci. USA 88:5202, 1991), lacks the carboxyl half of  
20 the third motif. Because removal of the carboxyl half of the first motif abolishes activity, it appears likely that the majority of the biological effectiveness of  $\eta$  can be attributed to the first two motifs. Although by different measures  $\eta$  is equally as active as  $\zeta$  in  
25 promoting antigen-mediated cytokine release (Bauer et al., Proc. Natl. Acad. Sci. USA 88:3842, 1991) or redirected cytotoxicity (see above),  $\eta$  is not phosphorylated in response to receptor stimulation (Bauer et al., supra, 1991). Thus either the presence of all three motifs is  
30 required for phosphorylation, or the third motif represents a favored substrate for an unidentified tyrosine kinase.

#### EXAMPLE IX

Cytolytic Signal Transduction by Human Fc Receptor

- 53 -

To evaluate the actions of different human Fc receptor subtypes, chimeric molecules were created in which the extracellular domain of the human CD4, CD5 or CD16 antigens were joined to the transmembrane and intracellular domains of the FcRII $\gamma$ A, B1, B2, and C subtypes (nomenclature of Ravetch and Kinet, Ann. Rev. Immunol. 9:457, 1991). Specifically, cDNA sequences corresponding to the transmembrane and cytoplasmic domains of the previously described FcRIIA, B1, and B2 isoforms were amplified from the preexisting clone PC23 or from a human tonsil cDNA library (constructed by standard techniques) using oligonucleotide primers:

CCC GGA TCC CAG CAT GGG CAG CTC TT (SEQ ID NO: 18; FcRIIA forward)

CGC GGG GCG GCC GCT TTA GTT ATT ACT GTT GAC ATG  
GTC GTT (SEQ ID NO: 19; FcRIIA reverse);

GCG GGG GGA TCC CAC TGT CCA AGC TCC CAG CTC TTC  
ACC G (SEQ ID NO: 20; FcRIIB1 and FcRIIB2 forward); and

GCG GGG GCG GCC GCC TAA ATA CGG TTC TGG TC (SEQ ID  
NO: 21; FcRIIB1 and FcRIIB2 reverse).

These primers contained cleavage sites for the enzymes BamHI and NotI, respectively, indented 6 residues from the 5' end. The NotI site was immediately followed by an antisense stop codon, either CTA or TTA. All primers contained 18 or more residues complementary to the 5' and 3' ends of the desired fragments. The cDNA fragment corresponding to the FcRII $\gamma$ C cytoplasmic domain, which differs from the IIA isoform in only one amino acid residue (L for P at residue 268) was generated by site directed mutagenesis by overlap PCR using primers of sequence:

TCA GAA AGA GAC AAC CTG AAG AAA CCA ACA A (SEQ ID  
NO: 22) and

TTG TTG GTT TCT TCA GGT TGT GTC TTT CTG A (SEQ ID  
NO: 23).

- 54 -

The PCR fragments were inserted into vaccinia virus expression vectors which contained the CD16 or CD4 extracellular domains respectively and subsequently inserted into wild type vaccinia by recombination at the thymidine kinase locus, using selection for cointegration of E coli gpt to facilitate identification of the desired recombinants. The identities of all isoforms (shown in Fig. 12) were confirmed by dideoxy sequencing.

Production of the chimeric receptor proteins was further confirmed by immunoprecipitation studies. Approximately  $10^7$  JRT3.T3.5 cells were infected for one hour in serum free IMDM medium with recombinant vaccinia at a multiplicity of infection of at least ten. Twelve hours post-infection, the cells were harvested and surface labeled with 0.5mCi  $^{125}\text{I}$  per  $10^7$  cells using the lactoperoxidase/glucose oxidase method (Clark and Einfeld, supra). The labeled cells were collected by centrifugation and lysed 1% NP-40, 0.1mM  $\text{MgCl}_2$ , 5mM KCl, 0.2M iodoacetamide and 1mM PMSF. Nuclei were removed by centrifugation, and CD16 fusion proteins immunoprecipitated with antibody 4G8 and anti-mouse IgG agarose. Samples were electrophoresed under reducing conditions. All immunoprecipitated chimeric receptor molecules were of the expected molecular masses.

To test the ability of the chimeric receptors to mediate an increase in cytoplasmic free calcium ion, the recombinant viruses were used to infect the TCR<sup>-</sup> mutant Jurkat cell line JRT3.T3.5 (as described herein) and cytoplasmic free calcium was measured in the cells (as described herein) following crosslinking of the receptor extracellular domains with monoclonal antibody 3G8 or Leu-3A (as described herein). These experiments revealed that the intracellular domains of FcR $\gamma$ II A and C were capable of mediating an increase in cytoplasmic free calcium ion after crosslinking of the extracellular

- 55 -

domains, whereas the intracellular domains of FcR $\gamma$ II B1 and B2 were inactive under comparable conditions (Fig. 13 A and 13B). The CD4, CD5 and CD16 hybrids of FcR $\gamma$ II A shared essentially equal capacity to promote the calcium response (Fig. 13 and data not shown). Other cell lines, from both monocytic and lymphocytic lineages, were capable of responding to the signal initiated by crosslinking of the extracellular domains (data not shown).

10 To explore the involvement of the different FcR $\gamma$ II intracellular domains in cytotoxicity, human cytotoxic T lymphocytes (CTL) were infected with vaccinia recombinants expressing CD16:FcR $\gamma$ II A, B1, B2 and C chimeras. The infected cells were then cocultured with  
15 <sup>51</sup>Cr-loaded hybridoma cells (i.e., 3G8 10-2 cells) which expressed cell surface antibody to CD16. In this assay CTLs bearing the CD16 chimera killed the hybridoma target cells (allowing release of free <sup>51</sup>Cr) if the CD16 extracellular domain of the chimera has been joined to an  
20 intracellular segment capable of activating the lymphocyte effector program; this cytotoxicity assay is described in detail below. Fig. 14A shows that CTL armed with CD16:FcR $\gamma$ IIA and C, but not FcR $\gamma$ II B1 or B2, are capable of lysing target cells expressing cell surface  
25 anti-CD16 antibody.

To eliminate the possibility that the specific cytotoxicity was in some way attributable to interaction with the CD16 moiety, cytotoxicity experiments were conducted in which the FcR $\gamma$ II intracellular domains were  
30 attached to a CD4 extracellular domain. In this case the target cells were HeLa cells expressing HIV envelope gp120/41 proteins (specifically, HeLa cells infected with the vaccinia vector vPE16 (available from the National Institute of Allergy and Infectious Disease AIDS  
35 Depository, Bethesda, MD). As in the CD16 system, target

- 56 -

cells expressing HIV envelope were susceptible to lysis by T cells expressing the CD4:FcR $\gamma$ II A chimera, but not FcR $\gamma$ II B1 or B2 (Fig. 14B).

The intracellular domains of FcR $\gamma$ II A and C share  
5 no appreciable sequence homology with any other protein, including the members of the extended FcR $\gamma$ /TCR $\zeta$  family. To define the sequence elements responsible for induction of cytolysis, 5' and 3' deletions of the intracellular domain coding sequences (described below and shown in  
10 Fig. 15A) were prepared and were evaluated for efficacy in calcium mobilization and cytolysis assays (as described herein). In the experiments in which the amino terminal portion of the intracellular domain was removed, the transmembrane domain of FcR $\gamma$ II was replaced with the  
15 transmembrane domain of the unrelated CD7 antigen to eliminate the possible contribution of interactions mediated by the membrane-spanning domain.

Figs. 15B and 15C show that removal of the 14  
carboxyl-terminal residues, including tyrosine 298,  
20 resulted in a complete loss of cytolytic capacity and a substantial reduction in calcium mobilization potential. Further deletion to just before tyrosine 282 gave an identical phenotype (Figs. 15B and 15C). Deletion from the N-terminus of the intracellular domain to residue 268  
25 had no substantial effect on either calcium profile or cytolytic potency, whereas deletion to residue 275 markedly impaired free calcium release but had little effect on cytolysis (Figs. 15D and 15E). Further  
deletion, to residue 282, gave FcR $\gamma$ II tails which lacked  
30 the ability to either mobilize calcium or trigger cytolysis (Figs. 15D and 15E). The 'active element' defined by these crude measures is relatively large (36 amino acids) and contains two tyrosines separated by 16 residues.

- 57 -

**EXAMPLE X**

Other intracellular and transmembrane signal transducing domains according to the invention may be derived from the T cell receptor proteins, CD3 delta and T3 gamma, and the B cell receptor proteins, mb1 and B29. The amino acid sequences of these proteins is shown in Fig. 16 (CD3 delta; SEQ ID NO: 24), Fig. 17 (T3 gamma; SEQ ID NO: 25), Fig. 18 (mb1; SEQ ID NO: 26) and Fig. 19 (B29; SEQ ID NO: 27). The portions of the sequences sufficient for cytolytic signal transduction (and therefore preferably included in a chimeric receptor of the invention) are shown in brackets. Chimeric receptors which include these protein domains are constructed and used in the therapeutic methods of the invention generally as described above.

**EXAMPLE XI****Experimental Methods****Vaccinia Infection and Radioimmunoprecipitation**

Approximately  $5 \times 10^6$  CV1 cells were infected for one hour in serum free DME medium with recombinant vaccinia at a multiplicity of infection (moi) of at least ten (titer measured on CV1 cells). The cells were placed in fresh medium after infection and labelled metabolically with  $200 \mu\text{Ci/ml}$   $^{35}\text{S}$ -methionine plus cysteine (Tran $^{35}\text{S}$ -label, ICN; Costo-Mesa, CA) in methionine and cysteine free DMEM (Gibco; Grand Island, NY) for six hours. The labelled cells were detached with PBS containing 1mM EDTA, collected by centrifugation, and lysed in 1% NP-40, 0.1% SDS, 0.15 M NaCl, 0.05M Tris pH 8.0, 5mM EDTA, and 1mM PMSF. Nuclei were removed by centrifugation, and CD4 proteins immunoprecipitated with OKT4 antibody and anti-mouse IgG agarose (Cappel, Durham, NC). Samples were electrophoresed through 8% polyacrylamide/SDS gels under non-reducing (NR) and reducing (R) conditions. Gels containing  $^{35}\text{S}$ -labelled

- 58 -

samples were impregnated with En<sup>3</sup>Hance (New England Nuclear, Boston, MA) prior to autoradiography.

Facilitated expression of the transmembrane form of CD16, CD16<sub>TM</sub>, was measured by comparing its expression in CV1  
5 cells singly infected with CD16<sub>TM</sub> with expression in cells coinfectd with viruses encoding CD16<sub>TM</sub> and  $\zeta$  or  $\gamma$  chimeras. After infection and incubation for six hours or more, cells were detached from plates with PBS, 1mM EDTA and the expression of CD16<sub>TM</sub> or the chimeras was  
10 measured by indirect immunofluorescence and flow cytometry.

#### Calcium Flux Assay

Jurkat subline E6 (Weiss et al., J. Immunol., 133:123-128 (1984)) cells were infected with recombinant  
15 vaccinia viruses for one hour in serum free IMDM at an moi of 10 and incubated for three to nine hours in IMDM, 10% FBS. Cells were collected by centrifugation and resuspended at  $3 \times 10^6$  cells/ml in complete medium containing 1mM Indo-1 acetomethoxyester (Grynkiewicz et  
20 al., J. Biol. Chem., 260:3340-3450 (1985)) (Molecular Probes) and incubated at 37°C for 45 minutes. The Indo-1 loaded cells were pelleted and resuspended at  $1 \times 10^6$ /ml in serum free IMDM and stored at room temperature in the dark. Cells were analyzed for free calcium ion by  
25 simultaneous measurement of the violet and blue fluorescence emission by flow cytometry (Rabinovitch et al., J. Immunol., 137:952-961 (1986)). To initiate calcium flux, either phycoerythrin (PE)-conjugated Leu-3A (anti-CD4) (Becton Dickinson, Lincoln Park, NJ) at 1  
30  $\mu$ g/ml was added to the cell suspension followed by 10 $\mu$ g/ml of unconjugated goat anti-mouse IgG at time 0 or unconjugated 3G8 (anti-CD16) monoclonal antibody was added to the cell suspension at 1  $\mu$ g/ml followed by 10  $\mu$ g/ml of PE-conjugated Fab<sub>2</sub>' goat anti-mouse IgG at time  
35 0. Histograms of the violet/blue emission ratio were



- 59 -

collected from the PE positive (infected) cell population, which typically represented 40-80% of all cells. The T cell antigen receptor response in uninfected cells was triggered by antibody OKT3, without crosslinking. For experiments involving CD16 chimeric receptors, samples showing baseline drift toward lower intracellular calcium (without antibody) were excluded from the analysis. Histogram data were subsequently analyzed by conversion of the binary data to ASCII using Write Hand Man (Cooper City, FL) software, followed by analysis with a collection of FORTRAN programs. The violet/blue emission ratio prior to the addition of the second antibody reagents was used to establish the normalized initial ratio, set equal to unity, and the resting threshold ratio, set so that 10% of the resting population would exceed threshold.

#### Cytolysis Assay

Human T cell line WH3, a CD8<sup>+</sup> CD4<sup>-</sup> HLA B44 restricted cytolytic line was maintained in IMDM, 10% human serum with 100 U/ml of IL-2 and was periodically stimulated either nonspecifically with irradiated (3000 rad) HLA-unmatched peripheral blood lymphocytes and 1µg/ml of phytohemagglutinin, or specifically, with irradiated B44-bearing mononuclear cells. After one day of nonspecific stimulation; the PHA was diluted to 0.5 µg/ml by addition of fresh medium, and after three days the medium was changed. Cells were grown for at least 10 days following stimulation before use in cytotoxicity assays. The cells were infected with recombinant vaccinia at a multiplicity of infection of at least 10 for one hour in serum free medium, followed by incubation in complete medium for three hours. Cells were harvested by centrifugation and resuspended at a density of  $1 \times 10^7$  cells/ml. 100µl were added to each well of a U-bottom microtiter plate containing 100 µl/well of complete

- 60 -

medium. Cells were diluted in two-fold serial steps. Two wells for each sample did not contain lymphocytes, to allow spontaneous chromium release and total chromium uptake to be measured. The target cells, from HeLa

5 subline S3, were infected in 6.0 or 10.0 cm plates at an approximate moi of 10 for one hour in serum free medium, followed by incubation in complete medium for three hours. They were then detached from the dishes with PBS, 1mM EDTA and counted. An aliquot of  $10^6$  target cells

10 (HeLa, Raji, or RJ2.2.5 cells for the CD4 chimeric receptor experiments and 3G8 10-2 cells; Shen et al., Mol. Immunol. 26:959 (1989) for the CD16 chimeric receptor experiments) was centrifuged and resuspended in 50  $\mu$ l of sterile  $^{51}\text{Cr}$ -sodium chromate (1mCi/ml, Dupont

15 Wilmington, DE) for one hour at 37°C with intermittent mixing, then washed three times with PBS. 100  $\mu$ l of labelled cells resuspended in medium at  $10^5$  cells/ml were added to each well. Raji and RJ2.2.5 target cells were labelled in the same manner as HeLa cells. The

20 microtiter plate was spun at 750 x g for 1 minute and incubated for 4 hours at 37°C. At the end of the incubation period, the cells in each well were resuspended by gentle pipetting, a sample removed to determine the total counts incorporated, and the

25 microtiter plate spun at 750 x g for 1 minute. 100 $\mu$ l aliquots of supernatant were removed and counted in a gamma ray scintillation counter. The percent killing was corrected for the fraction of infected target cells (usually 50-90%) measured by flow cytometry. For

30 infected effector cells the effector:target ratio was corrected for the percent of cells infected (usually 20-50% for the CD4 chimeric receptor experiments and >70% for the CD16 chimeric receptor experiments).

- 61 -

**In Vitro Mutagenesis of the  $\zeta$  Sequence**

To create point mutations in amino acid residues 11 and or 15 of the  $\zeta$  sequence, synthetic oligonucleotide primers extending from the BamHI site upstream of the  $\zeta$  transmembrane domain, and converting native  $\zeta$  residue 11 from Cys to Gly (C11G) or residue 15 from Asp to Gly (D15G) or both (C11G/D15G) were prepared and used in PCR reactions to generate mutated fragments which were reinserted into the wild type CD4: $\zeta$  constructs.

- 10 To create  $\zeta$  deletions,  $\zeta$  cDNA sequences were amplified by PCR using synthetic oligonucleotide primers designed to create a stop codon (UAG) after residues 50, 59, or 65. The primers contained the cleavage site for the enzyme NotI indented five or six residues from the 5' end, usually in a sequence of the form CGC GGG CGG CCG CTA (SEQ ID NO: 11), where the last three residues correspond to the stop anticodon. The NotI and stop anticodon sequences were followed by 18 or more residues complementary to the desired 3' end of the fragment.
- 20 The resulting chimeras were designated CD16: $\zeta$ Y51\*, CD16: $\zeta$ E60\* and CD16: $\zeta$ D66\* respectively. The BamHI site upstream of the transmembrane domain and the NotI site were used to generate fragments that were reintroduced into the wild type CD16: $\zeta$  construct. Monomeric  $\zeta$
- 25 chimeras were created by liberating the  $\zeta$  transmembrane and membrane proximal intracellular sequences by BamHI and SacI digestion of the Asp<sup>-</sup> and Cys<sup>-</sup> CD4: $\zeta$  construct described above and inserting the fragment into the CD16: $\zeta$ E60\* and CD16: $\zeta$ D66\* construct respectively.
- 30 CD16:7: $\zeta$ (48-65) and CD16:7: $\zeta$ (48-59) tripartite chimera construction.

To prepare the construct CD16: $\zeta$ D66\*, the  $\zeta$  cDNA sequence corresponding to the transmembrane domain and the 17 following residues of the cytoplasmic domain was

35 replaced by corresponding transmembrane and cytoplasmic

- 62 -

domain obtained from the CD5 and CD7 cDNA. The CD5 and CD7 fragments were generated by a PCR reaction using forward oligonucleotides including a BamHI restriction cleavage site and corresponding to the region just upstream of the transmembrane domain of CD5 and CD7 respectively and the following reverse oligonucleotides overlapping the CD5 and CD7 sequences respectively and the  $\zeta$  sequence which contained the SacI restriction cleavage site.

10 CD5: $\zeta$ : CGC GGG CTC GTT ATA GAG CTG GTT CTG GCG  
CTG CTT CTT CTG (SEQ ID NO: 12)

CD7: $\zeta$ : CGC GGG GAG CTC GTT ATA GAG CTG GTT TGC  
CGC CGA ATT CTT ATC CCG (SEQ ID NO: 13).

The CD5 and CD7 PCR products were digested with BamHI and SacI and ligated to BamHI and SacI digested CD16: $\zeta$ E60\* and replacing the  $\zeta$  sequence from BamHI to SacI by the CD7 fragment. To make the constructs CD16:CD5 and CD16:CD7, CD5 and CD7 fragments were obtained by PCR using an oligonucleotide containing a NotI restriction cleavage site and encoding a stop codon (UAA) after the residue Gln416 and Ala193 of CD5 and CD7 respectively. The CD5 and CD7 PCR fragment were digested with BamHI and NotI and inserted in the CD16: $\zeta$ Asp66\* construct.

25 **In Vitro Mutagenesis of the N-terminal Residues within the  $\zeta$  Cytolytic Signal-Transducing Motif**

Synthetic oligonucleotide primers extending from the SacI site inside the  $\zeta$  motif and converting native residue 48 from Asn to Ser (N48S), residue 50 from Leu to Ser (L50S) and residue 51 from Tyr to Phe (Y51F) were synthesized and used in a PCR reaction to generate fragments that were reintroduced into the wild type CD16:7: $\zeta$ (48-65) construct.

**In Vitro Mutagenesis of C-terminal Residues within the  $\zeta$  Cytolytic Signal-Transducing Motif**

- 63 -

Synthetic oligonucleotide primers extending from the NotI site 3' to the stop codon and converting native residue 60 from Glu to Gln (E60Q), residue 61 from Glu to Gln (E61Q), residue 62 from Tyr to Phe or Ser (Y62F or Y62S) and residue 63 from Asp to Asn (D63N) were synthesized and used in PCR to generate fragments that were subcloned into the wild type CD16:5D66\* construct from the BamHI site to the NotI site.

CD16:7:5(33-65), CD16:7:5(71-104), CD16:7:5(104-137)

#### 10 Chimera Constructions

A CD7 transmembrane fragment bearing MluI and NotI sites at the junction between the transmembrane and intracellular domains was obtained by PCR using an oligonucleotide with the following sequence: CGC GGG GCG GCC ACG CGT CCT CGC CAG CAC ACA (SEQ ID NO: 14). The resulting PCR fragment was digested with BamHI and NotI and reinserted into the CD16:7:5(48-65) construct. 5 fragments encoding residues 33 to 65, 71 to 104, and 104 to 137 were obtained by PCR reaction using pairs of primers containing MluI sites at the 5' end of the forward primers and stop codons followed by NotI sites at the 5' end of the reverse primers. In each case the restriction sites were indented six residues from the 5' terminus of the primer to insure restriction enzyme cleavage.

5 33: CGC GGG ACG CGT TTC AGC CGT CCT CGC CAG CAC ACA (SEQ ID NO: 15);

5 71: CGC GGG ACG CGT GAC CCT GAG ATG GGG GGA AAG (SEQ ID NO: 16); and

30 5 104: CGC GGG ACG CGT ATT GGG ATG AAA GGC GAG CGC (SEQ ID NO: 17).

#### Construction of FcR $\gamma$ IIA Deletion Mutants

Carboxyl terminal FcRIIA deletion mutants were constructed by PCR in the same fashion as for the full length constructs, converting the sequences encoding

- 64 -

tyrosine at positions 282 and 298 into stop codons (TAA). The N-terminal deletions were generated by amplifying fragments encoding successively less of the intracellular domain by PCR, using oligonucleotides which allowed the  
5 resulting fragments to be inserted between MluI and NotI restriction sites into a previously constructed expression plasmid encoding the CD16 extracellular domain fused to the CD7 transmembrane domain, the latter terminating in a MluI site and the junction between the  
10 transmembrane and the intracellular domain.

#### OTHER EMBODIMENTS

The examples described above demonstrate that aggregation of  $\zeta$ ,  $\eta$ , or  $\gamma$  chimeras suffices to initiate the cytolytic effector cell response in T cells. The  
15 known range of expression of  $\zeta$ ,  $\eta$ , and  $\gamma$ , which includes T lymphocytes, natural killer cells, basophilic granulocytes, macrophages and mast cells, suggests that conserved sequence motifs may interact with a sensory apparatus common to cells of hematopoietic origin and  
20 that an important component of host defense in the immune system may be mediated by receptor aggregation events.

The potency of the cytolytic response and the absence of a response to target cells bearing MHC class II receptors demonstrates that chimeras based on  $\zeta$ ,  $\eta$ , or  
25  $\gamma$  form the basis for a genetic intervention for AIDS through adoptive immunotherapy. The broad distribution of endogenous  $\zeta$  and  $\gamma$  and evidence that Fc receptors associated with  $\gamma$  mediate cytotoxicity in different cells types (Fanger et al., Immunol. Today, 10:92-99 (1989))  
30 allows a variety of cells to be considered for this purpose. For example, neutrophilic granulocytes, which have a very short lifespan ( $\approx$  4h) in circulation and are intensely cytolytic, are attractive target cells for expression of the chimeras. Infection of neutrophils  
35 with HIV is not likely to result in virus release, and

- 65 -

the abundance of these cells (the most prevalent of the leukocytes) should facilitate host defense. Another attractive possibility for host cells are mature T cells, a population presently accessible to retroviral engineering (Rosenberg, S.A. Sci. Am., 262:62-69 (1990)). With the aid of recombinant IL-2, T cell populations can be expanded in culture with relative ease, and the expanded populations typically have a limited lifespan when reinfused (Rosenberg et al., N. Engl. J. Med., 323:570-578 (1990)).

Under the appropriate conditions, HIV recognition by cells expressing CD4 chimeras should also provide mitogenic stimuli, allowing the possibility that the armed cell population could respond dynamically to the viral burden. Although we have focused here on the behavior of the fusion proteins in cytolytic T lymphocytes, expression of the chimeras in helper lymphocytes might provide an HIV-mobilized source of cytokines which could counteract the collapse of the helper cell subset in AIDS. Recent description of several schemes for engineering resistance to infection at steps other than virus penetration (Friedman et al., Nature, 335:452-454 (1988); Green et al., Cell, 58:215-223 (1989); Malim et al., Cell, 58:205-214 (1989); Trono et al., Cell, 59:113-120 (1989); Buonocore et al., Nature, 345:625-628 (1990)) suggests that cells bearing CD4 chimeras could be designed to thwart virus production by expression of appropriate agents having an intracellular site of action.

The ability to transmit signals to T lymphocytes through autonomous chimeras also provides the ability for the regulation of retrovirally engineered lymphocytes in vivo. Crosslinking stimuli, mediated for example by specific IgM antibodies engineered to remove complement-binding domains, may allow such lymphocytes to increase

- 66 -

in number in situ, while treatment with similar specific IgG antibodies (for example recognizing an amino acid variation engineered into the chimeric chain) could selectively deplete the engineered population.

- 5 Additionally, anti-CD4 IgM antibodies do not require additional crosslinking to mobilize calcium in Jurkat cells expressing CD4: $\zeta$  chimeras (data not shown). The ability to regulate cell populations without recourse to repeated extracorporeal amplification may substantially  
10 extend the range and efficacy of current uses proposed for genetically engineered T cells.

To create other chimeras consisting of  $\zeta$ ,  $\eta$  or  $\gamma$  intracellular sequences, cDNA or genomic sequences encoding an extracellular domain of the receptor can be  
15 endowed with a restriction site introduced at a location just preceding the transmembrane domain of choice. The extracellular domain fragment terminating in the restriction site can then be joined to  $\zeta$ ,  $\eta$ , or  $\gamma$  sequences. Typical extracellular domains may be derived  
20 from receptors which recognize complement, carbohydrates, viral proteins, bacteria, protozoan or metazoan parasites, or proteins induced by them. Similarly, ligands or receptors expressed by pathogens or tumor cells can be attached to  $\zeta$ ,  $\eta$ , or  $\gamma$  sequences, to direct  
25 immune responses against cells bearing receptors recognizing those ligands.

While the invention has been described in connection with specific embodiments thereof, it will be understood that it is capable of further modifications  
30 and this application is intended to cover variations, uses, or adaptations of the invention and including such departures from the present disclosure as come within the art to which the invention pertains and as may be applied to the essential features hereinbefore set forth as  
35 follows in the scope of the appended claims.



- 67 -

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

- (i) APPLICANTS: The General Hospital Corporation
- (ii) TITLE OF INVENTION: Redirection of Cellular Immunity by Receptor  
Chimeras
- (iii) NUMBER OF SEQUENCES: 27
- (iv) CORRESPONDENCE ADDRESS:
  - (A) ADDRESSEE: Fish & Richardson
  - (B) STREET: 225 Franklin Street
  - (C) CITY: Boston
  - (D) STATE: MA
  - (E) COUNTRY: USA
  - (F) ZIP: 02110-2804
- (v) COMPUTER READABLE FORM:
  - (A) MEDIUM TYPE: 3.5" Diskette, 1.44 Mb
  - (B) COMPUTER: IBM PS/2 Model 50Z or 55SX
  - (C) OPERATING SYSTEM: IBM P.C. DOS (Version 3.30)
  - (D) SOFTWARE: Wordperfect (Version 5.0)
- (vi) CURRENT APPLICATION DATA:
  - (A) APPLICATION NUMBER:
  - (B) FILING DATE:
  - (C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:
  - (A) APPLICATION NUMBER: 07/665,961
  - (B) FILING DATE: March 7, 1991
  - (C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
  - (A) NAME: Clark, Paul T.
  - (B) REGISTRATION NUMBER: 30,162
  - (C) REFERENCE/DOCKET NUMBER: 00786/119002
- (ix) TELECOMMUNICATION INFORMATION:
  - (A) TELEPHONE: (617) 542-5070
  - (B) TELEFAX: (617) 542-8906
  - (C) TELEX: 200154

## (2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 1728 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)

- 68 -

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

ATGAACCGGG GAGTCCCTTT TAGGCACTTG CTTCTGGTGC TGCAACTGGC	50
GCTCCTCCCA GCAGCCACTC AGGGAAACAA AGTGGTGCTG GGCAAAAAAG	100
GGGATACAGT GGAAC TGACC TGTACAGCTT CCCAGAAGAA GAGCATACAA	150
TTCCACTGGA AAAACTCCAA CCAGATAAAG ATTCTGGGAA ATCAGGGGCTC	200
CTTCTTAACT AAAGGTCCAT CCAAGCTGAA TGATCGCGCT GACTCAAGAA	250
GAAGCCTTTG GGACCAAGGA AACTTCCCCC TGATCATCAA GAATCTTAAG	300
ATAGAAGACT CAGATACTTA CATCTGTGAA GTGGAGGACC AGAAGGAGGA	350
GGTGCAATTG CTAGTGTTG GATTGACTGC CAACTCTGAC ACCCACCTGC	400
TTCAGGGGCA GAGCCTGACC CTGACCTTGG AGAGCCCCC TGGTAGTAGC	450
CCCTCAGTGC AATGTAGGAG TCCAAGGGGT AAAAACATAC AGGGGGGGAA	500
GACCCTCTCC GTGTCTCAGC TGGAGCTCCA GGATAGTGGC ACCTGGACAT	550
GCACTGTCTT GCAGAACCAG AAGAAGGTGG AGTTCAAAAT AGACATCGTG	600
GTGCTAGCTT TCCAGAAGGC CTCCAGCATA GTCTATAAGA AAGAGGGGGA	650
ACAGGTGGAG TTCTCCTTCC CACTCGCCTT TACAGTTGAA AAGCTGACGG	700
GCAGTGGCGA GCTGTGGTGG CAGGCGGAGA GGGCTTCCTC CTCCAAGTCT	750
TGGATCACCT TTGACCTGAA GAACAAGGAA GTGTCTGTAA AACGGGTTAC	800
CCAGGACCCT AAGCTCCAGA TGGGCAAGAA GCTCCCGCTC CACCTCACCC	850
TGCCCCAGGC CTTGCCTCAG TATGCTGGCT CTGGAAACCT CACCCTGGCC	900
CTTGAAGCGA AAACAGGAAA GTTGCATCAG GAAGTGAACC TGGTGGTGAT	950
GAGAGCCACT CAGCTCCAGA AAAATTTGAC CTGTGAGGTG TGGGGACCCA	1000
CCTCCCCTAA GCTGATGCTG AGCTTGAAAC TGGAGAACAA GGAGGCAAAG	1050
GTCTCGAAGC GGGAGAAGCC GGTGTGGGTG CTGAACCTG AGGCGGGGAT	1100
GTGGCAGTGT CTGCTGAGTG ACTCGGGACA GGTCTGTCTG GAATCCAACA	1150
TCAAGGTTCT GCCCACATGG TCCACCCCGG TGCACGCGGA TCCCAAACCTC	1200
TGCTACTTGC TAGATGGAAT CCTCTTCATC TACGGAGTCA TCATCACAGC	1250
CCTGTACCTG AGAGCAAAAT TCAGCAGGAG TGCAGAGACT GCTGCCAACC	1300

- 69 -

TGCAGGACCC CAACCAGCTC TACAATGAGC TCAATCTAGG GCGAAGAGAG	1350
GAATATGACG TCTTGGAGAA GAAGCGGGCT CGGGATCCAG AGATGGGAGG	1400
CAAACAGCAG AGGAGGAGGA ACCCCCAGGA AGGCGTATAC AATGCACTGC	1450
AGAAAGACAA GATGCCAGAA GCCTACAGTG AGATCGGCAC AAAAGGCGAG	1500
AGGCGGAGAG GCAAGGGGCA CGATGGCCTT TACCAGGACA GCCACTTCCA	1550
AGCAGTGCAG TTCGGGAACA GAAGAGAGAG AGAAGGTTCA GAACTCACAA	1600
GGACCCTTGG GTTAAGAGCC CGCCCCAAAG GTGAAAGCAC CCAGCAGAGT	1650
AGCCAATCCT GTGCCAGCGT CTTCAGCATC CCCACTCTGT GGAGTCCATG	1700
GCCACCCAGT AGCAGCTCCC AGCTCTAA	1728

## (2) INFORMATION FOR SEQ ID NO:2:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1389 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA (genomic)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

ATGAACCGGG GAGTCCCTTT TAGGCACTTG CTTCTGGTGC TGCAACTGGC	50
GCTCCTCCCA GCAGCCACTC AGGGAAACAA AGTGGTGCTG GGCAAAAAAG	100
GGGATACAGT GGAAGTGACC TGTACAGCTT CCCAGAAGAA GAGCATACAA	150
TTCCAAGTGA AAAACTCCAA CCAGATAAAG ATTCTGGGAA ATCAGGGGCTC	200
CTTCTTAACT AAAGGTCCAT CCAAGCTGAA TGATCGCGCT GACTCAAGAA	250
GAAGCCTTTG GGACCAAGGA AACTTCCCCC TGATCATCAA GAATCTTAAG	300
ATAGAAGACT CAGATACTTA CATCTGTGAA GTGGAGGACC AGAAGGAGGA	350
GGTGCAATTG CTAGTGTTTG GATTGACTGC CAACTCTGAC ACCCACCTGC	400
TTCAGGGGCA GAGCCTGACC CTGACCTTGG AGAGCCCCC TGGTAGTAGC	450
CCCTCAGTGC AATGTAGGAG TCCAAGGGGT AAAAACATAC AGGGGGGGAA	500
GACCTCTCC GTGTCTCAGC TGGAGCTCCA GGATAGTGGC ACCTGGACAT	550
GCACTGTCTT GCAGAACCAG AAGAAGGTGG AGTTCAAAT AGACATCGTG	600

- 70 -

GTGCTAGCTT TCCAGAAGGC CTCCAGCATA GTCTATAAGA AAGAGGGGGA	650
ACAGGTGGAG TTCTCCTTCC CACTCGCCTT TACAGTTGAA AAGCTGACGG	700
GCAGTGGCGA GCTGTGGTGG CAGGCGGAGA GGGCTTCCTC CTCCAAGTCT	750
TGGATCACCT TTGACCTGAA GAACAAGGAA GTGTCTGTAA AACGGGTTAC	800
CCAGGACCCT AAGCTCCAGA TGGGCAAGAA GCTCCCGCTC CACCTCACCC	850
TGCCCCAGGC CTTGCCTCAG TATGCTGGCT CTGGAAACCT CACCCTGGCC	900
CTTGAAGCGA AAACAGGAAA GTTGCAATCAG GAAGTGAACC TGGTGGTGTAT	950
GAGAGCCACT CAGCTCCAGA AAAATTTGAC CTGTGAGGTG TGGGGACCCA	1000
CCTCCCCATA GCTGATGCTG AGCTTGAAAC TGGAGAACAA GGAGGCAAAG	1050
GTCTCGAAGC GGGAGAAGCC GGTGTGGGTG CTGAACCCTG AGGCGGGGAT	1100
GTGGCAGTGT CTGCTGAGTG ACTCGGGACA GGTCTGCTG GAATCCAACA	1150
TCAAGGTTCT GCCACATGG TCCACCCCGG TGCACGCGGA TCCGCAGCTC	1200
TGCTATATCC TGGATGCCAT CCTGTTTTTG TATGGTATTG TCCTTACCCT	1250
GCTCTACTGT CGACTCAAGA TCCAGGTCCG AAAGGCAGAC ATAGCCAGCC	1300
GTGAGAAATC AGATGCTGTC TACACGGGCC TGAACACCCG GAACCAGGAG	1350
ACATATGAGA CTCTGAAACA TGAGAAACCA CCCCAATAG	1389

## (2) INFORMATION FOR SEQ ID NO:3:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1599 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA (genomic)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

ATGAACCGGG GAGTCCCTTT TAGGCACTTG CTTCTGGTGC TGCAACTGGC	50
GCTCCTCCCA GCAGCCACTC AGGGAAACAA AGTGGTGCTG GGCAAAAAG	100
GGGATACAGT GGAAGTGACC TGTACAGCTT CCCAGAAGAA GAGCATACAA	150
TTCCACTGGA AAAACTCCAA CCAGATAAAG ATTCTGGGAA ATCAGGGCTC	200
CTTCTTAACT AAAGGTCCAT CCAAGCTGAA TGATCGCGCT GACTCAAGAA	250

- 71 -

GAAGCCTTTG GGACCAAGGA AACTTCCCCC TGATCATCAA GAATCTTAAG	300
ATAGAAGACT CAGATACTTA CATCTGTGAA GTGGAGGACC AGAAGGAGGA	350
GGTGCAATTG CTAGTGTTTCG GATTGACTGC CAACTCTGAC ACCCACCTGC	400
TTCAGGGGCA GAGCCTGACC CTGACCTTGG AGAGCCCCC TGGTAGTAGC	450
CCCTCAGTGC AATGTAGGAG TCCAAGGGGT AAAAACATAC AGGGGGGGAA	500
GACCCTCTCC GTGTCTCAGC TGGAGCTCCA GGATAGTGGC ACCTGGACAT	550
GCACTGTCTT GCAGAACCAG AAGAAGGTGG AGTTCAAAAT AGACATCGTG	600
GTGCTAGCTT TCCAGAAGGC CTCCAGCATA GTCTATAAGA AAGAGGGGGA	650
ACAGGTGGAG TTCTCCTTCC CACTCGCCTT TACAGTTGAA AAGCTGACGG	700
GCAGTGGCGA GCTGTGGTGG CAGGCGGAGA GGGCTTCCTC CTCCAAGTCT	750
TGGATCACCT TTGACCTGAA GAACAAGGAA GTGTCTGTAA AACGGGTTAC	800
CCAGGACCCT AAGCTCCAGA TGGGCAAGAA GCTCCCGCTC CACCTCACCC	850
TGCCCCAGGC CTTGCCTCAG TATGCTGGCT CTGGAAACCT CACCCTGGCC	900
CTTGAAGCGA AAACAGGAAA GTTGCATCAG GAAGTGAACC TGGTGGTGAT	950
GAGAGCCACT CAGCTCCAGA AAAATTTGAC CTGTGAGGTG TGGGGACCCA	1000
CCTCCCCTAA GCTGATGCTG AGCTTGAAAC TGGAGAACAA GGAGGCAAAG	1050
GTCTCGAAGC GGGAGAAGCC GGTGTGGGTG CTGAACCTG AGGCGGGGAT	1100
GTGGCAGTGT CTGCTGAGTG ACTCGGGACA GGTCTGCTG GAATCCAACA	1150
TCAAGGTTCT GCCCACATGG TCCACCCCGG TGCACGCGGA TCCCAAATC	1200
TGCTACCTGC TGGATGGAAT CCTCTTCATC TATGGTGTCA TTCTCACTGC	1250
CTGTTCCTG AGAGTGAAGT TCAGCAGGAG CGCAGAGCCC CCCGCGTACC	1300
AGCAGGGCCA GAACCAGCTC TATAACGAGC TCAATCTAGG ACGAAGAGAG	1350
GAGTACGATG TTTTGGACAA GAGACGTGGC CGGGACCCTG AGATGGGGGG	1400
AAAGCCGAGA AGGAAGAACC CTCAGGAAGG CCTGTACAAT GAACTGCAGA	1450
AAGATAAGAT GGCAGGAGCC TACAGTGAGA TTGGGATGAA AGGCGAGCGC	1500
CGGAGGGGCA AGGGGCACGA TGGCCTTTAC CAGGGTCTCA GTACAGCCAC	1550
CAAGGACACC TACGACGCCC TTCACATGCA GGCCCTGCCC CCTCGCTAA	1599

(2) INFORMATION FOR SEQ ID NO:4:

- 72 -

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 575 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Met	Asn	Arg	Gly	Val	Pro	Phe	Arg	His	Leu	Leu	Leu	Val	Leu	Gln	Leu	1	5	10	15
Ala	Leu	Leu	Pro	Ala	Ala	Thr	Gln	Gly	Asn	Lys	Val	Val	Leu	Gly	Lys	20	25	30	
Lys	Gly	Asp	Thr	Val	Glu	Leu	Thr	Cys	Thr	Ala	Ser	Gln	Lys	Lys	Ser	35	40	45	
Ile	Gln	Phe	His	Trp	Lys	Asn	Ser	Asn	Gln	Ile	Lys	Ile	Leu	Gly	Asn	50	55	60	
Gln	Gly	Ser	Phe	Leu	Thr	Lys	Gly	Pro	Ser	Lys	Leu	Asn	Asp	Arg	Ala	65	70	75	80
Asp	Ser	Arg	Arg	Ser	Leu	Trp	Asp	Gln	Gly	Asn	Phe	Pro	Leu	Ile	Ile	85	90	95	
Lys	Asn	Leu	Lys	Ile	Glu	Asp	Ser	Asp	Thr	Tyr	Ile	Cys	Glu	Val	Glu	100	105	110	
Asp	Gln	Lys	Glu	Glu	Val	Gln	Leu	Val	Phe	Gly	Leu	Thr	Ala	Asn		115	120	125	
Ser	Asp	Thr	His	Leu	Leu	Gln	Gly	Gln	Ser	Leu	Thr	Leu	Thr	Leu	Glu	130	135	140	
Ser	Pro	Pro	Gly	Ser	Ser	Pro	Ser	Val	Gln	Cys	Arg	Ser	Pro	Arg	Gly	145	150	155	160
Lys	Asn	Ile	Gln	Gly	Gly	Lys	Thr	Leu	Ser	Val	Ser	Gln	Leu	Glu	Leu	165	170	175	
Gln	Asp	Ser	Gly	Thr	Trp	Thr	Cys	Thr	Val	Leu	Gln	Asn	Gln	Lys	Lys	180	185	190	
Val	Glu	Phe	Lys	Ile	Asp	Ile	Val	Val	Leu	Ala	Phe	Gln	Lys	Ala	Ser	195	200	205	
Ser	Ile	Val	Tyr	Lys	Lys	Glu	Gly	Glu	Gln	Val	Glu	Phe	Ser	Phe	Pro	210	215	220	
Leu	Ala	Phe	Thr	Val	Glu	Lys	Leu	Thr	Gly	Ser	Gly	Glu	Leu	Trp	Trp	225	230	235	240
Gln	Ala	Glu	Arg	Ala	Ser	Ser	Ser	Lys	Ser	Trp	Ile	Thr	Phe	Asp	Leu	245	250	255	
Lys	Asn	Lys	Glu	Val	Ser	Val	Lys	Arg	Val	Thr	Gln	Asp	Pro	Lys	Leu	260	265	270	
Gln	Met	Gly	Lys	Lys	Leu	Pro	Leu	His	Leu	Thr	Leu	Pro	Gln	Ala	Leu	275	280	285	
Pro	Gln	Tyr	Ala	Gly	Ser	Gly	Asn	Leu	Thr	Leu	Ala	Leu	Glu	Ala	Lys	290	295	300	
Thr	Gly	Lys	Leu	His	Gln	Glu	Val	Asn	Leu	Val	Val	Met	Arg	Ala	Thr	305	310	315	320
Gln	Leu	Gln	Lys	Asn	Leu	Thr	Cys	Glu	Val	Trp	Gly	Pro	Thr	Ser	Pro	325	330	335	
Lys	Leu	Met	Leu	Ser	L u	Lys	Leu	Glu	Asn	Lys	Glu	Ala	Lys	Val	Ser	340	345	350	
Lys	Arg	lu	Lys	Pro	Val	Trp	Val	Leu	Asn	Pro	Glu	Ala	Gly	Met	Trp	355	360	365	

- 73 -

```

In Cys Leu Leu Ser Asp Ser Gly Gln Val Leu Leu Glu Ser Asn Ile
370                      375                      380
Lys Val Leu Pro Thr Trp Ser Thr Pro Val His Ala Asp Pro Lys Leu
385                      390                      395                      400
Cys Tyr Leu Leu Asp Gly Ile Leu Phe Ile Tyr Gly Val Ile Ile Thr
405                      410                      415
Ala Leu Tyr Leu Arg Ala Lys Phe Ser Arg Ser Ala Glu Thr Ala Ala
420                      425                      430
Asn Leu Gln Asp Pro Asn Gln Leu Tyr Asn Glu Leu Asn Leu Gly Arg
435                      440                      445
Arg Glu Glu Tyr Asp Val Leu Glu Lys Lys Arg Ala Arg Asp Pro Glu
450                      455                      460
Met Gly Gly Lys Gln Gln Arg Arg Arg Asn Pro Gln Glu Gly Val Tyr
465                      470                      475                      480
Asn Ala Leu Gln Lys Asp Lys Met Pro Glu Ala Tyr Ser Glu Ile Gly
485                      490                      495
Thr Lys Gly Glu Arg Arg Arg Gly Lys Gly His Asp Gly Leu Tyr Gln
500                      505                      510
Asp Ser His Phe Gln Ala Val Gln Phe Gly Asn Arg Arg Glu Arg Glu
515                      520                      525
Gly Ser Glu Leu Thr Arg Thr Leu Gly Leu Arg Ala Arg Pro Lys Gly
530                      535                      540
Glu Ser Thr Gln Gln Ser Ser Gln Ser Cys Ala Ser Val Phe Ser Ile
555                      560                      565                      570
Pro Thr Leu Trp Ser Pro Trp Pro Pro Ser Ser Ser Ser Gln Leu
565                      570                      575

```

## (2) INFORMATION FOR SEQ ID NO:5:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 462 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

```

Met Asn Arg Gly Val Pro Phe Arg His Leu Leu Leu Val Leu Gln Leu
1           5           10           15
Ala Leu Leu Pro Ala Ala Thr Gln Gly Asn Lys Val Val Leu Gly Lys
20           25           30
Lys Gly Asp Thr Val Glu Leu Thr Cys Thr Ala Ser Gln Lys Lys Ser
35           40           45
Ile Gln Phe His Trp Lys Asn Ser Asn Gln Ile Lys Ile Leu Gly Asn
50           55           60
Gln Gly Ser Phe Leu Thr Lys Gly Pro Ser Lys Leu Asn Asp Arg Ala
65           70           75           80
Asp Ser Arg Arg Ser Leu Trp Asp Gln Gly Asn Phe Pro Leu Ile Ile
85           90           95
Lys Asn Leu Lys Ile Glu Asp Ser Asp Thr Tyr Il Cys Glu Val lu
100          105          110
Asp Gln Lys Glu Glu Val Gln Leu Leu Val Phe Gly Leu Thr Ala Asn
115          120          125
Ser Asp Thr His Leu Leu Gln Gly Gln Ser Leu Thr Leu Thr Leu lu

```

- 74 -

130		135		140
Ser Pro Pro Gly Ser Ser Pro Ser Val Gln Cys Arg Ser Pro Arg Gly				
145		150		155
Lys Asn Ile Gln Gly Gly Lys Thr Leu Ser Val Ser Gln Leu Glu Leu				160
	165		170	175
Gln Asp Ser Gly Thr Trp Thr Cys Thr Val Leu Gln Asn Gln Lys Lys				
	180		185	190
Val Glu Phe Lys Ile Asp Ile Val Val Leu Ala Phe Gln Lys Ala Ser				
	195		200	205
Ser Ile Val Tyr Lys Lys Glu Gly Glu Gln Val Glu Phe Ser Phe Pro				
	210		215	220
Leu Ala Phe Thr Val Glu Lys Leu Thr Gly Ser Gly Glu Leu Trp Trp				
225		230		235
Gln Ala Glu Arg Ala Ser Ser Ser Lys Ser Trp Ile Thr Phe Asp Leu				240
	245		250	255
Lys Asn Lys Glu Val Ser Val Lys Arg Val Thr Gln Asp Pro Lys Leu				
	260		265	270
Gln Met Gly Lys Lys Leu Pro Leu His Leu Thr Leu Pro Gln Ala Leu				
	275		280	285
Pro Gln Tyr Ala Gly Ser Gly Asn Leu Thr Leu Ala Leu Glu Ala Lys				
	290		295	300
Thr Gly Lys Leu His Gln Glu Val Asn Leu Val Val Met Arg Ala Thr				
305		310		315
Gln Leu Gln Lys Asn Leu Thr Cys Glu Val Trp Gly Pro Thr Ser Pro				
	325		330	335
Lys Leu Met Leu Ser Leu Lys Leu Glu Asn Lys Glu Ala Lys Val Ser				
	340		345	350
Lys Arg Glu Lys Pro Val Trp Val Leu Asn Pro Glu Ala Gly Met Trp				
	355		360	365
Gln Cys Leu Leu Ser Asp Ser Gly Gln Val Leu Leu Glu Ser Asn Ile				
	370		375	380
Lys Val Leu Pro Thr Trp Ser Thr Pro Val His Ala Asp Pro Gln Leu				
385		390		395
Cys Tyr Ile Leu Asp Ala Ile Leu Phe Leu Tyr Gly Ile Val Leu Thr				
	405		410	415
Leu Leu Tyr Cys Arg Leu Lys Ile Gln Val Arg Lys Ala Asp Ile Ala				
	420		425	430
Ser Arg Glu Lys Ser Asp Ala Val Tyr Thr Gly Leu Asn Thr Arg Asn				
	435		440	445
Gln Glu Thr Tyr Glu Thr Leu Lys His Glu Lys Pro Pro Gln				
450		455		460
				462

## (2) INFORMATION FOR SEQ ID NO:6:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 532 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Met Asn Arg Gly Val Pro Ph	Arg His Leu Leu Leu Val Leu Gln Leu
1	5 10 15



- 75 -

Ala	Leu	Leu	Pro	Ala	Ala	Thr	Gln	Gly	Asn	Lys	Val	Val	Leu	Gly	Lys		
			20					25					30				
Lys	Gly	Asp	Thr	Val	Glu	Leu	Thr	Cys	Thr	Ala	Ser	Gln	Lys	Lys	Ser		
		35					40					45					
Ile	Gln	Phe	His	Trp	Lys	Asn	Ser	Asn	Gln	Ile	Lys	Ile	Leu	Gly	Asn		
	50					55					60						
Gln	Gly	Ser	Phe	Leu	Thr	Lys	Gly	Pro	Ser	Lys	Leu	Asn	Asp	Arg	Ala		
	65				70					75					80		
Asp	Ser	Arg	Arg	Ser	Leu	Trp	Asp	Gln	Gly	Asn	Phe	Pro	Leu	Ile	Ile		
				85					90						95		
Lys	Asn	Leu	Lys	Ile	Glu	Asp	Ser	Asp	Thr	Tyr	Ile	Cys	Glu	Val	Glu		
			100					105					110				
Asp	Gln	Lys	Glu	Glu	Val	Gln	Leu	Leu	Val	Phe	Gly	Leu	Thr	Ala	Asn		
		115					120					125					
Ser	Asp	Thr	His	Leu	Leu	Gln	Gly	Gln	Ser	Leu	Thr	Leu	Thr	Leu	Glu		
	130					135					140						
Ser	Pro	Pro	Gly	Ser	Ser	Pro	Ser	Val	Gln	Cys	Arg	Ser	Pro	Arg	Gly		
	145				150					155					160		
Lys	Asn	Ile	Gln	Gly	Gly	Lys	Thr	Leu	Ser	Val	Ser	Gln	Leu	Glu	Leu		
			165						170					175			
Gln	Asp	Ser	Gly	Thr	Trp	Thr	Cys	Thr	Val	Leu	Gln	Asn	Gln	Lys	Lys		
			180					185					190				
Val	Glu	Phe	Lys	Ile	Asp	Ile	Val	Val	Leu	Ala	Phe	Gln	Lys	Ala	Ser		
		195					200					205					
Ser	Ile	Val	Tyr	Lys	Lys	Glu	Gly	Glu	Gln	Val	Glu	Phe	Ser	Phe	Pro		
	210					215					220						
Leu	Ala	Phe	Thr	Val	Glu	Lys	Leu	Thr	Gly	Ser	Gly	Glu	Leu	Trp	Trp		
	225				230					235					240		
Gln	Ala	Glu	Arg	Ala	Ser	Ser	Ser	Lys	Ser	Trp	Ile	Thr	Phe	Asp	Leu		
				245					250					255			
Lys	Asn	Lys	Glu	Val	Ser	Val	Lys	Arg	Val	Thr	Gln	Asp	Pro	Lys	Leu		
			260					265					270				
Gln	Met	Gly	Lys	Lys	Leu	Pro	Leu	His	Leu	Thr	Leu	Pro	Gln	Ala	Leu		
		275				280						285					
Pro	Gln	Tyr	Ala	Gly	Ser	Gly	Asn	Leu	Thr	Leu	Ala	Leu	Glu	Ala	Lys		
	290					295					300						
Thr	Gly	Lys	Leu	His	Gln	Glu	Val	Asn	Leu	Val	Val	Met	Arg	Ala	Thr		
	305				310					315					320		
Gln	Leu	Gln	Lys	Asn	Leu	Thr	Cys	Glu	Val	Trp	Gly	Pro	Thr	Ser	Pro		
				325					330					335			
Lys	Leu	Met	Leu	Ser	Leu	Lys	Leu	Glu	Asn	Lys	Glu	Ala	Lys	Val	Ser		
		340						345					350				
Lys	Arg	Glu	Lys	Pro	Val	Trp	Val	Leu	Asn	Pro	Glu	Ala	Gly	Met	Trp		
		355					360					365					
Gln	Cys	Leu	Leu	Ser	Asp	Ser	Gly	Gln	Val	Leu	Leu	Glu	Ser	Asn	Ile		
	370					375						380					
Lys	Val	Leu	Pro	Thr	Trp	Ser	Thr	Pro	Val	His	Ala	Asp	Pro	Lys	Leu		
	385				390					395					400		
Cys	Tyr	Leu	Leu	Asp	Gly	Ile	Leu	Phe	Ile	Tyr	Gly	Val	Ile	Leu	Thr		
				405					410					415			
Ala	Leu	Phe	Leu	Arg	Val	Lys	Phe	Ser	Arg	Ser	Ala	lu	Pr	Pro	Ala		
			420					425					430				
Tyr	Gln	ln	Gly	ln	Asn	Gln	Leu	Tyr	Asn	Glu	Leu	Asn	Leu	ly	Arg		
		435					440					445					
Arg	lu	Glu	Tyr	Asp	Val	Leu	Asp	Lys	Arg	Arg	Gly	Arg	Asp	Pro	lu		
	450					455						460					

- 76 -

Met Gly Gly Lys Pro Arg Arg Lys Asn Pro Gln Glu Gly Leu Tyr Asn  
 465 470 475 480  
 Lu Leu Gln Lys Asp Lys Met Ala Glu Ala Tyr Ser Glu Ile Gly Met  
 485 490 495  
 Lys Gly Glu Arg Arg Arg Gly Lys Gly His Asp Gly Leu Tyr Gln Gly  
 500 505 510  
 Leu Ser Thr Ala Thr Lys Asp Thr Tyr Asp Ala Leu His Met Gln Ala  
 515 520 525  
 Leu Pro Pro Arg  
 530

## (2) INFORMATION FOR SEQ ID NO:7:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 33 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

CGCGGGGTGA CCGTGCCCTC CAGCAGCTTG GGC

33

## (2) INFORMATION FOR SEQ ID NO:8:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 50 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

CGCGGGGATC CGTCGTCCAG AGCCCGTCCA GCTCCCCGTC CTGGGCCTCA

50

## (2) INFORMATION FOR SEQ ID NO:9:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 33 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: nucleic acid

- 77 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

CGCGGGGCGGC CGCGACGCCG GCCAAGACAG CAC

33

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 33 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

CGCGTTGACG AGCAGCCAGT TGGGCAGCAG CAG

33

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 15 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

CGCGGGGCGGC CGCTA

15

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 42 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

CGCGGGGCTCG TTATAGAGCT GGTTCCTGGCG CTGCTTCTTC TG

42

(2) INFORMATION FOR SEQ ID NO:13:

- 78 -

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 48 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

CGCGGGGAGC TCGTTATAGA GCTGGTTTGC CGCCGAATTC TTATCCCG

48

## (2) INFORMATION FOR SEQ ID NO:14:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 33 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

CGCGGGGCGG CCACGCGTCC TCGCCAGCAC ACA

33

## (2) INFORMATION FOR SEQ ID NO:15:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 36 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

CGCGGGGACGC GTTTCAGCCG TCCTCGCCAG CACACA

36

## (2) INFORMATION FOR SEQ ID NO:16:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 33 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOP LOGY: linear

(ii) MOLECULE TYPE: nucleic acid

- 79 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

CGCGGGACGC GTGACCCTGA GATGGGGGCA AAG

33

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 33 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

CGCGGGACGC GTATTGGGAT GAAAGGCGAG CGC

33

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

CCCGGATCCC AGCATGGGCA GCTCTT

26

(2) INFORMATION FOR SEQ ID NO:19:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 42 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

CGCGGGGCGG CCGCTTTAGT TATTACTGTT GACATGGTCG TT

42

(2) INFORMATION FOR SEQ ID NO:20:

(i) SEQUENCE CHARACTERISTICS:

- 80 -

- (A) LENGTH: 30 bases
- (B) TYPE: nucl ic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

GCGGGGGGAT CCCACTGTCC AAGCTCCCAG

30

(2) INFORMATION FOR SEQ ID NO:21:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 32 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

GCGGGGGCGG CCGCCTAAAT ACGGTTCTGG TC

32

(2) INFORMATION FOR SEQ ID NO:22:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 31 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

TCAGAAAGAG ACAACCTGAA GAAACCAACA A

31

(2) INFORMATION FOR SEQ ID NO:23:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 31 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: nucleic acid

- 81 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

TTGTTGGTTT CTTCAGGTTG TGTCTTTCTG A

31

(2) INFORMATION FOR SEQ ID NO:24:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 171 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: amino acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

```

Met Glu His Ser Thr Phe Leu Ser Gly Leu Val Leu Ala Thr Leu Leu
      5              10              15
Ser Gln Val Ser Pro Phe Lys Ile Pro Ile Glu Glu Leu Glu Asp Arg
      20              25              30
Val Phe Val Asn Cys Asn Thr Ser Ile Thr Trp Val Glu Gly Thr Val
      35              40              45
Gly Thr Leu Leu Ser Asp Ile Thr Arg Leu Asp Leu Gly Lys Arg Ile
      50              55              60
Leu Asp Pro Arg Gly Ile Tyr Arg Cys Asn Gly Thr Asp Ile Tyr Lys
      65              70              75              80
Asp Lys Glu Ser Thr Val Gln Val His Tyr Arg Met Cys Gln Ser Cys
      85              90              95
Val Glu Leu Asp Pro Ala Thr Val Ala Gly Ile Ile Val Thr Asp Val
      100             105             110
Ala Ile Thr Leu Leu Leu Ala Leu Gly Val Phe Cys Phe Ala Gly His
      115             120             125
Glu Thr Gly Arg Leu Ser Gly Ala Ala Asp Thr Gln Ala Leu Leu Arg
      130             135             140
Asn Asp Gln Val Tyr Gln Pro Leu Arg Asp Arg Asp Asp Ala Gln Tyr
      145             150             155             160
Ser His Leu Gly Gly Asn Trp Ala Arg Asn Lys
      165             170

```

(2) INFORMATION FOR SEQ ID NO:25:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 182 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: amino acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

```

Met Glu Gln Gly Lys Gly Leu Ala Val Leu Ile Leu Ala Ile Ile Leu
      5              10              15

```

- 82 -

```

Leu  In Gly Thr Leu Ala Gln Ser Ile Lys Gly Asn His Leu Val Lys
      20                      25                      30
Val Tyr Asp Tyr  In Glu Asp Gly Ser Val Leu Leu Thr Cys Asp Ala
      35                      40                      45
Glu Ala Lys Asn Ile Thr Trp Phe Lys Asp Gly Lys Met Ile Gly Phe
      50                      55                      60
Leu Thr Glu Asp Lys Lys Lys Trp Asn Leu Gly Ser Asn Ala Lys Asp
      65                      70                      75                      80
Pro Arg Gly Met Tyr Gln Cys Lys Gly Ser Gln Asn Lys Ser Lys Pro
      85                      90                      95
Leu Gln Val Tyr Tyr Arg Met Cys Gln Asn Cys Ile Glu Leu Asn Ala
      100                     105                     110
Ala Thr Ile Ser Gly Phe Leu Phe Ala Glu Ile Val Ser Ile Phe Val
      115                     120                     125
Leu Ala Val Gly Val Tyr Phe Ile Ala Gly Gln Asp Gly Val Arg Gln
      130                     135                     140
Ser Arg Ala Ser Asp Lys Gln Thr Leu Leu Pro Asn Asp Gln Leu Tyr
      145                     150                     155                     160
Gln Pro Leu Lys Asp Arg Glu Asp Asp Gln Tyr Ser His Leu Gln Gly
      165                     170                     175
Asn Gln Leu Arg Arg Asn
      180

```

## (2) INFORMATION FOR SEQ ID NO:26:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 220 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: amino acids

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

```

Met Pro Gly Gly Leu Glu Ala Leu Arg Ala Leu Pro Leu Leu Leu Phe
      5                      10                      15
Leu Ser Tyr Ala Cys Leu Gly Pro Gly Cys Gln Ala Leu Arg Val Glu
      20                      25                      30
Gly Gly Pro Pro Ser Leu Thr Val Asn Leu Gly Glu Glu Ala Arg Leu
      35                      40                      45
Thr Cys Glu Asn Asn Gly Arg Asn Pro Asn Ile Thr Trp Trp Phe Ser
      50                      55                      60
Leu Gln Ser Asn Ile Thr Trp Pro Pro Val Pro Leu Gly Pro Gly Gln
      65                      70                      75                      80
Gly Thr Thr Gly Gln Leu Phe Phe Pro Glu Val Asn Lys Asn Thr Gly
      85                      90                      95
Ala Cys Thr Gly Cys Gln Val Ile Glu Asn Asn Ile Leu Lys Arg Ser
      100                     105                     110
Cys Gly Thr Tyr Leu Arg Val Arg Asn Pro Val Pro Arg Pro Phe Leu
      115                     120                     125
Asp Met Gly Glu  ly Thr Lys Asn Arg Ile Ile Thr Ala Glu Gly Ile
      130                     135                     140
Ile Leu Leu Phe Cys Ala Val Val Pro Gly Thr Leu Leu Leu Phe Arg
      145                     150                     155                     160

```



- 83 -

```

Lys Arg Trp Gln Asn Glu Lys Phe Gly Val Asp Met Pro Asp Asp Tyr
                      165                      170                      175
Glu Asp Glu Asn Leu Tyr Glu Gly Leu Asn Leu Asp Asp Cys Ser Met
                      180                      185                      190
Tyr Glu Asp Ile Ser Arg Gly Leu Gln Gly Thr Tyr Gln Asp Val Gly
                      195                      200                      205
Asn Leu His Ile Gly Asp Ala Gln Leu Glu Lys Pro
                      210                      215                      220

```

## (2) INFORMATION FOR SEQ ID NO:27:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 228 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: amino acid

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

```

Met Ala Thr Leu Val Leu Ser Ser Met Pro Cys His Trp Leu Leu Phe
                      5                      10                      15
Leu Leu Leu Leu Phe Ser Gly Glu Pro Val Pro Ala Met Thr Ser Ser
                      20                      25                      30
Asp Leu Pro Leu Asn Phe Gln Gly Ser Pro Cys Ser Gln Ile Trp Gln
                      35                      40                      45
His Pro Arg Phe Ala Ala Lys Lys Arg Ser Ser Met Val Lys Phe His
                      50                      55                      60
Cys Tyr Thr Asn His Ser Gly Ala Leu Thr Trp Phe Arg Lys Arg Gly
                      65                      70                      75                      80
Ser Gln Gln Pro Gln Glu Leu Val Ser Glu Glu Gly Arg Ile Val Gln
                      85                      90                      95
Thr Gln Asn Gly Ser Val Tyr Thr Leu Thr Ile Gln Asn Ile Gln Tyr
                      100                     105                     110
Glu Asp Asn Gly Ile Tyr Phe Cys Lys Gln Lys Cys Asp Ser Ala Asn
                      115                     120                     125
His Asn Val Thr Asp Ser Cys Gly Thr Glu Leu Leu Val Leu Gly Phe
                      130                     135                     140
Ser Thr Leu Asp Gln Leu Lys Arg Arg Asn Thr Leu Lys Asp Gly Ile
                      145                     150                     155                     160
Ile Leu Ile Gln Thr Leu Leu Ile Ile Leu Phe Ile Ile Val Pro Ile
                      165                     170                     175
Phe Leu Leu Leu Asp Lys Asp Asp Gly Lys Ala Gly Met Glu Glu Asp
                      180                     185                     190
His Thr Tyr Glu Gly Leu Asn Ile Asp Gln Thr Ala Thr Tyr Glu Asp
                      195                     200                     205
Ile Val Thr Leu Arg Thr Gly Glu Val Lys Trp Ser Val Gly Glu His
                      210                     215                     220
Pro Gly  In Glu
225

```

- 84 -

Claims

1           1. A method of directing a cellular response to an  
2 infective agent, to a cell infected with said agent, to a  
3 tumor or cancerous cell, or to an autoimmune-generated cell in  
4 a mammal, said method comprising administering to said mammal  
5 an effective amount of therapeutic cells, said cells being  
6 capable of specifically recognizing and destroying said agent  
7 or said cell.

1           2. The method of claim 1, wherein said cellular response  
2 is MHC-independent.

1           3. The method of claim 1, wherein said therapeutic cells  
2 express a membrane-bound, proteinaceous chimeric receptor,  
3 comprising (a) an extracellular portion which is capable of  
4 specifically recognizing and binding said agent or said cell,  
5 and (b) an intracellular or transmembrane portion which is  
6 capable of signalling said therapeutic cell to destroy a  
7 receptor-bound agent or a receptor-bound cell.

1           4. The method of claim 3, wherein said intracellular  
2 portion or said transmembrane portion is a signal-transducing  
3 portion of a T cell receptor protein, an Fc receptor protein,  
4 or a B cell receptor protein, or a functional derivative  
5 thereof.

1           5. The method of claim 4, wherein said receptor protein  
2 is  $\epsilon$ ,  $\eta$ ,  $\gamma$ , CD3 delta, T3 gamma, mb1, or B29.

1           6. The method of claim 5, wherein said chimeric receptor  
2 comprises:

- 3           (a) amino acids 421-532 of SEQ ID NO: 6;  
4           (b) amino acids 423-455, 438-455, 461-494 or 494-528 of  
5 SEQ ID NO:6;  
6           (c) amino acids 400-420 of SEQ ID NO:6;  
7           (d) amino acids 421-575 of SEQ ID NO: 4;

- 85 -

8 (e) amino acids 423-455, 438-455, 461-494 or 494-528 of  
9 SEQ ID NO:4;

10 (f) amino acids 400-420 of SEQ ID NO: 4;

11 (g) amino acids 421-462 of SEQ ID NO: 5;

12 (h) amino acids 402-419 of SEQ ID NO:5;

13 (i) amino acids Tyr282-Tyr298 inclusive of Fig. 15A;

14 (j) amino acids 132-171 of Fig. 16 (SEQ ID NO: 24);

15 (k) amino acids 140-182 of Fig. 17 (SEQ ID NO: 25);

16 (l) amino acids 162-220 of Fig. 18 (SEQ ID NO: 26);

17 (m) amino acids 183-228 of Fig. 19 (SEQ ID NO: 27).

1 7. The method of claim 1, wherein said therapeutic cells  
2 are selected from the group consisting of:

3 (a) T lymphocytes;

4 (b) cytotoxic T lymphocytes;

5 (c) natural killer cells;

6 (d) neutrophils;

7 (e) granulocytes;

8 (f) macrophages;

9 (g) mast cells;

10 (h) HeLa cells; and

11 (i) embryonic stem cells (ES).

1 8. The method of claim 1, wherein said infective agent  
2 is an immunodeficiency virus.

1 9. The method of claim 8, wherein said extracellular  
2 portion comprises an HIV envelope-binding portion of CD4, or a  
3 functional HIV envelope-binding derivative thereof.

1 10. The method of claim 9, wherein said cytotoxic T  
2 lymphocyte is a CD8<sup>+</sup> cytotoxic T lymphocytes.

1 11. A cell which expresses a proteinaceous membrane-  
2 bound chimeric receptor, said receptor comprising (a) an  
3 extracellular portion which is capable of specifically  
4 recognizing and binding an infective agent, a cell infected

- 86 -

5 with an infective agent, a tumor or cancerous cell, or an  
6 autoimmune-generated cell, and (b) an intracellular portion or  
7 transmembrane portion derived from a T cell receptor, an Fc  
8 receptor, or a B cell receptor which is capable of signalling  
9 said cell to destroy a receptor-bound agent or receptor-bound  
10 cell.

1 12. The cell of claim 11, wherein said binding is MHC-  
2 independent.

1 13. The cell of claim 11, wherein said receptor protein  
2 is  $\zeta$ ,  $\eta$ ,  $\gamma$ , CD3 delta, T3 gamma, mb1, or B29.

1 14. The cell of claim 13, wherein said chimeric receptor  
2 comprises amino acids:.

3 (a) amino acids 421-532 of SEQ ID NO: 6;

4 (b) amino acids 423-455, 438-455, 461-494 or 494-528 of  
5 SEQ ID NO:6;

6 (c) amino acids 400-420 of SEQ ID NO:6;

7 (d) amino acids 421-575 of SEQ ID NO: 4;

8 (e) amino acids 423-455, 438-455, 461-494 or 494-528 of  
9 SEQ ID NO:4;

10 (f) amino acids 400-420 of SEQ ID NO: 4;

11 (g) amino acids 421-462 of SEQ ID NO: 5;

12 (h) amino acids 402-419 of SEQ ID NO:5;

13 (i) amino acids Tyr282-Tyr298 inclusive of Fig. 15A;

14 (j) amino acids 132-171 of Fig. 16 (SEQ ID NO: 24);

15 (k) amino acids 140-182 of Fig. 17 (SEQ ID NO: 25);

16 (l) amino acids 162-220 of Fig. 18 (SEQ ID NO: 26);

17 (m) amino acids 183-228 of Fig. 19 (SEQ ID NO: 27).

1 15. The cell of claim 11, wherein said therapeutic cells  
2 are selected from the group consisting of:

3 (a) T lymphocytes;

4 (b) cytotoxic T lymphocytes;

5 (c) natural killer cells;

6 (d) neutrophils;

- 87 -

- 7 (e) granulocytes;
- 8 (f) macrophages;
- 9 (g) mast cells;
- 10 (h) HeLa cells; and
- 11 (i) embryonic stem cells (ES).

1 16. The cell of claim 11, wherein said extracellular  
2 portion comprises an HIV envelope-binding portion of CD4, or a  
3 functional HIV envelope-binding derivative thereof.

1 17. The cell of claim 16, wherein said cytotoxic T  
2 lymphocyte is a CD8<sup>+</sup> cytotoxic T lymphocytes.


1 18. The cell of claim 11, wherein said extracellular  
2 portion is an immunoglobulin molecule, or an antigen-binding  
3 fragment thereof.

1 19. DNA encoding a chimeric receptor of claim 11.

1 20. A vector comprising the chimeric receptor DNA of  
2 claim 19.

1 21. A method of treating an immunodeficiency virus  
2 infection, comprising administering the host cell of claim 16  
3 to a mammal.

1 22. A substantially pure antibody which specifically  
2 recognizes and binds a chimeric receptor of claim 11.



1 / 22




FIG. 1a




FIG. 1b


SUBSTITUTE SHEET







2 / 22

**FIG. 1c**




**FIG. 2**







3 / 22

**FIG. 3a**



**FIG. 3b**





4 / 22

FIG. 4a







FIG. 4b



SUBSTITUTE SHEET



5 / 22

**FIG. 4c****FIG. 4d****SUBSTITUTE SHEET**





6 / 22

FIG. 5a






FIG. 5b




SUBSTITUTE SHEET



7 / 22

**FIG. 5c****SUBSTITUTE SHEET**



8 / 22

FIG. 6a





FIG. 6b



SUBSTITUTE SHEET



9 / 22

FIG. 7a






FIG. 7b




SUBSTITUTE SHEET









11 / 22

FIG. 9a





FIG. 9b



SUBSTITUTE SHEET



12 / 22

FIG. 9c





FIG. 9d



SUBSTITUTE SHEET



13 / 22

FIG. 10a






FIG. 10b



SUBSTITUTE SHEET





14 / 22

FIG. 10c






FIG. 10f



SUBSTITUTE SHEET



15 / 22

FIG. 10d






FIG. 10e



SUBSTITUTE SHEET



16 / 22

FIG. 11a







FIG. 11b



SUBSTITUTE SHEET



17 / 22



**FIG. 12**





18 / 22

FIG. 13a





FIG. 13b



SUBSTITUTE SHEET



19 / 22

FIG. 14a






FIG. 14b



SUBSTITUTE SHEET



20 / 22

FIG. 15a




FIG. 15b





FIG. 15c



SUBSTITUTE SHEET



21 / 22

FIG. 15d





FIG. 15e



SUBSTITUTE SHEET





22 / 22

FIG.16 (Seq. ID No: 24)

1 MEHSTFLSGL VLATLLSQVS PFKIPIEELE DRVFNVCNTS ITWVEGTVGT  
 51 LLSDITRLDL GKRILDPRGI YRCNGTDIYK DKESTVQVHY RMCQSCVELD  
 101 PATVAGIIVT DVIATLLLLAL GVFCFAGHET GRLSGAADTQ ALLRNDQVYQ  
 151 PLRDRDDAQY SHLGGNWARN K\*

FIG.17 (Seq ID NO: 25)

1 MEQGKGLAVL ILAIILLQGT LAQSIKGNHL VKVYDYQEDG SVLLTCDAEA  
 51 KNITWFKDGK MIGFLTEDKK KWNLGSSNAKD PRGMYQCKGS QNKSPLQVY  
 101 YRMCQNCIEL NAATISGFLF AEIVSIFVLA VGVYFIAGQD GVRQSRASDK  
 151 QTLLPNDQLY QPLKDREDDQ YSHLQGNQLR RN\*

FIG.18 (Seq ID No: 26)

1 MPGGLEALRA LPLLLFLSYA CLGPGCQALR VEGGPPSLTV NLGEEARLTC  
 51 ENNGRNPNTIT WWFSLQSNIT WPPVPLGPGQ GTTGQLFFPE VNKNTGACTG  
 101 CQVIENNILK RSCGTYLRVR NPVPRPFLDM GEGTKNRIIT AEGIILLFCA  
 151 VVPGTLLLFRR KRWQNEKFGV DMPDDYEDEN LYEGLNLDDC SMYEDISRGL  
 201 QGTYQDVGNL HIGDAQLEKP \*

FIG.19 (Seq ID No: 27)

1 MATLVLSMP CHWLLFLLLL FSSEPVPAMT SSDLPLNFQG SPCSQIWQHP  
 51 RFAAKKRSSM VKFHCYTNHS GALTWFRKRG SQQPQELVSE EGRIVQTQNG  
 101 SVYTLTIQNI QYEDNGIYFC KQKCDSANHN VTDSCGTELL VLGFSTLDQL  
 151 KRRNTLKDGI ILIQTLIIIL FIIVPIFLLL DKDDGKAGME EDHTYEGLNI  
 201 DQTATYEDIV TLRTGEVKWS VGEHPGQE\*



# INTERNATIONAL SEARCH REPORT

International Application No. PCT/US92/01785

<b>I. CLASSIFICATION OF SUBJECT MATTER</b> (if several classification symbols apply, indicate all) <sup>3</sup> According to International Patent Classification (IPC) or to both National Classification and IPC IPC (5): A61K 37/12; C07K 3/00, 13/00, 15/00, 17/00 US CL : 530/387						
<b>II. FIELDS SEARCHED</b> <div style="text-align: center; border: 1px solid black; padding: 2px; margin: 5px 0;">Minimum Documentation Searched<sup>4</sup></div> <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 20%; padding: 5px;">Classification System</td> <td style="padding: 5px;">Classification Symbols</td> </tr> <tr> <td style="padding: 5px;">U.S.</td> <td style="padding: 5px;">530/387</td> </tr> </table> <div style="text-align: center; border: 1px solid black; padding: 2px; margin: 5px 0;">Documentation Searched other than Minimum Documentation to the extent that such Documents are included in the Fields Searched<sup>5</sup></div> <p style="padding: 5px;">APS, Dialog, Intelligenetics</p>			Classification System	Classification Symbols	U.S.	530/387
Classification System	Classification Symbols					
U.S.	530/387					
<b>III. DOCUMENTS CONSIDERED TO BE RELEVANT</b> <sup>14</sup>						
Category <sup>*</sup>	Citation of Document, <sup>16</sup> with indication, where appropriate, of the relevant passages <sup>17</sup>	Relevant to Claim No. <sup>18</sup>				
X,P	Cell, Vol. 64, issued 08 March 1991, C.Romeo & B. Seed, "Cellular immunity to HIV activated by CD4 fused to T cell or Fc receptor polypeptides", pages 1037-1046, see entire document.	1-22				
Y	Proc. Natl. Acad. Sci. USA, Vol. 85, issued December 1988, A.M. Weissman et al., "Molecular cloning and chromosomal localization of the human T-cell receptor zeta chain: Distinction from the molecular CD3 complex", pages 9709-9713, see entire document.	1-22				
Y	J. Biol. Chem., Vol 264, No. 35, issued 15 December 1989, S.A. Carr et al., "Protein and carbohydrate structural analysis of a recombinant soluble CD4 receptor by mass spectrometry", pages 21286-21295, see entire document.	1-22				
Y	Proc. Natl. Acad. Sci. USA, Vol. 83, issued May 1986, P. van den Elsen et al., "Exon/intron organization of the genes coding for the $\delta$ chains of the human and murine T-cell receptor/T3 complex", pages 2944-2948, see entire document.	1-22				
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p><sup>*</sup> Special categories of cited documents:<sup>15</sup></p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</p> <p>"&amp;" document member of the same patent family</p> </div> </div>						
<b>IV. CERTIFICATION</b>						
Date of the Actual Completion of the International Search <sup>2</sup>	Date of Mailing of this International Search Report <sup>2</sup>					
03 June 1992	12 JUN 1992					
International Searching Authority <sup>1</sup>	Signature of Authorized Officer <sup>20</sup>					
ISA/US	 KAREN COCHRANE CARLSON, PH.D.					

## FURTHER INFORMATION CONTINUED FROM THE SECOND SHEET

Y	DNA and Cell Biology, Vol. 9, No. 5, issued 1990, G. Zettlmeissl et al., "Expression and characterization of human CD4: Immunoglobulin fusion proteins", pages 347-353, see entire document.	1-22
Y	EMBO J, Vol. 6, No. 10, issued 1987, A. Tunnacliffe et al., "Physical linkage of three CD3 genes on human chromosome 11", pages 2953-2957, see entire document.	1-22

V. ☐ OBSERVATIONS WHERE CERTAIN CLAIMS WERE FOUND UNSEARCHABLE<sup>1</sup>

1. ☐ Claim numbers , because they relate to subject matter (1) not required to be searched by this Authority, namely:
2. ☐ Claim numbers , because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out (1), specifically:
3. ☐ Claim numbers , because they are dependent claims not drafted in accordance with the second and third sentences of PCT Rule 8.4(a).

VI. ☐ OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING<sup>2</sup>

This International Searching Authority found multiple inventions in this international application as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims of the international application.
2. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims of the international application for which fees were paid, specifically claims:
3. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claim numbers:
4. ☐ As all searchable claims could be searched without effort justifying an additional fee, the International Search Authority did not invite payment of any additional fee.

Remark on protest

- ☐ The additional search fees were accompanied by applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category*	Citation of Document, <sup>16</sup> with indication, where appropriate, of the relevant passages <sup>17</sup>	Relevant to Claim No. <sup>18</sup>
Y	EMBO J, vol. 5, No. 8, issued 1986, G.W. Krissansen et al., "Primary structure of the T3 gamma subunit of the T3/T cell antigen receptor complex deduced from cDNA sequences: evolution of the T3 gamma and delta subunits", pages 1799-1808, see entire document.	1-22
Y	EMBO J, Vol. 7, No. 11, issued 1988, N. Sakaguchi et al., "B lymphocyte lineage-restricted expression of MB-1, a gene with CD3-like structural properties", pages 3457-3464, see entire document.	1-22
Y	Proc. Natl. Acad. Sci. USA, Vol. 85, issued September 1988, G.G. Hermanson et al., "B29: A member of the immunoglobulin gene superfamily exclusively expressed on B-lineage cells", pages 6890-6894, see entire document.	1-22
Y	Proc. Natl. Acad. Sci. USA, Vol. 87, issued May 1990, Y. Jin et al., "Molecular cloning of the CD3 eta subunit identifies a CD3 zeta-related product in thymus-derived cells", pages 3319-3323, see entire document.	1-22
Y	J. Biol. Chem., Vol. 265, No. 11, issued 15 April 1990, H. Kuster et al., "Characterization and expression of the gene for the human Fc receptor gamma subunit", pages 6448-6452, see entire document.	1-22

**THIS PAGE BLANK (USPTO)**

**THIS PAGE BLANK (USPTO)**

**THIS PAGE BLANK (USPTO)**